

Baxter d -permutations and other pattern avoiding classes


Nicolas Bonichon and Pierre-Jean Morel

Journal of Integer Sequences Vol. 25: Article 22.8.3, 2022

Permutations and diagrams

A **permutation** $\sigma = \sigma(1), \dots, \sigma(n)$ is a bijection from $[n] := \{1, 2, \dots, n\}$ to itself. $P_\sigma := \{(i, \sigma(i)), i \in [n]\}$ the **diagram of σ** .


$$\sigma = 413526$$



Permutations and diagrams

A **permutation** $\sigma = \sigma(1), \dots, \sigma(n)$ is a bijection from $[n] := \{1, 2, \dots, n\}$ to itself. $P_\sigma := \{(i, \sigma(i)), i \in [n]\}$ the **diagram of σ** .

$$\sigma = 413526$$




P_σ :

A **diagram** (of size n) is a point set on $[n] \times [n]$ with exactly **1 point per row and 1 point per column**.

Permutations and diagrams

A **permutation** $\sigma = \sigma(1), \dots, \sigma(n)$ is a bijection from $[n] := \{1, 2, \dots, n\}$ to itself. $P_\sigma := \{(i, \sigma(i)), i \in [n]\}$ the **diagram of σ** .

$$\sigma = 413526$$



P_σ :

A **diagram** (of size n) is a point set on $[n] \times [n]$ with exactly **1 point per row and 1 point per column**.

S_n the set of permutations of size n .

$$|S_n| = n!$$

The Starting Question


- Permutations \Leftrightarrow Diagrams

The Starting Question

- Permutations \Leftrightarrow Diagrams
- Diagrams are 2D objects

The Starting Question

- Permutations \Leftrightarrow Diagrams
- Diagrams are 2D objects
- What could be "3D" diagrams?
- \Rightarrow "3D" Permutations?




d -Diagrams and d -Permutations

A **3-diagram** of size n is a point set on $[n]^3$ such that

d -Diagrams and d -Permutations

A **3-diagram** of size n is a point set on $[n]^3$ such that each plane of the grid (orthogonal to x, y or z) contains exactly 1 point.




Points:

- (1,2,6)
- (2,5,5)
- (3,3,4)
- (4,1,3)
- (5,4,2)
- (6,6,1)

A d -diagram is a point set of size n on $[n]^d$ such that each hyperplane orthogonal $x_i = j$ with $i \in [d]$ and $j \in [n]$ contains exactly 1 point.

d -Diagrams and d -Permutations

A **3-permutation** $\sigma := (\sigma_y, \sigma_z)$ is a couple of permutations.
 $S_n^2 := \{\text{3-permutations of size } n\}$. $|S_n^2| = n!^2$.



Points:

(1,2,6)

(2,5,5)

(3,3,4)

(253146, 654321)


(4,1,3)

(5,4,2)

(6,6,1)

A **d -permutation** of size n , $\sigma := (\sigma_1, \dots, \sigma_{d-1})$ is a sequence of $d-1$ permutations of size n . $S_n^{d-1} := \{d\text{-permutations of size } n\}$. $|S_n^{d-1}| = n!^{d-1}$.

Projections



$$\text{proj}_{xy}(\sigma) = \sigma_y$$

$$\text{proj}_{xz}(\sigma) = \sigma_z$$


$$\text{proj}_{yz}(\sigma) = \sigma_z \sigma_y^{-1}$$

$$\text{proj}_{yz}((253146, 654321)) = 364251$$

$$\text{proj}_{yx}(\sigma) = \sigma_y^{-1}$$

Let $\bar{\sigma} := (\text{Id}_n, \sigma_1, \dots, \sigma_{d-1})$, the **projection** on i of d -permutation σ is the d' -permutation $\text{proj}_i(\sigma) := \bar{\sigma}_{i_2} \bar{\sigma}_{i_1}^{-1}, \bar{\sigma}_{i_3} \bar{\sigma}_{i_1}^{-1}, \dots, \bar{\sigma}_{i_{d'}} \bar{\sigma}_{i_1}^{-1}$. d' is the **dimension** of the projection.

Projections



$$\text{proj}_{xy}(\sigma) = \sigma_y$$

$$\text{proj}_{xz}(\sigma) = \sigma_z$$


$$\text{proj}_{yz}(\sigma) = \sigma_z \sigma_y^{-1}$$

$$\text{proj}_{yz}((253146, 654321)) = 364251$$

$$\text{proj}_{yx}(\sigma) = \sigma_y^{-1} \text{ (indirect)}$$

Let $\bar{\sigma} := (\text{Id}_n, \sigma_1, \dots, \sigma_{d-1})$, the **projection** on i of d -permutation σ is the d' -permutation $\text{proj}_i(\sigma) := \bar{\sigma}_{i_2} \bar{\sigma}_{i_1}^{-1}, \bar{\sigma}_{i_3} \bar{\sigma}_{i_1}^{-1}, \dots, \bar{\sigma}_{i_{d'}} \bar{\sigma}_{i_1}^{-1}$. d' is the **dimension** of the projection.


Pattern (classic)



$\sigma = 413526$ contains the pattern $\pi = 213$.

A permutation σ **contains** a permutation (or a **pattern**) $\pi = \pi(1), \dots, \pi(k) \in S_k$ if there exist indices $c_1 < \dots < c_k$ such that $\sigma(c_1) \dots \sigma(c_k)$ is order-isomorphic to π . The set of points of indices c_1, \dots, c_k is an **occurrence** of the π .


Pattern (classic)



$\sigma = 413526$ contains the pattern $\pi = 213$.
Positions 1,5,6 is an **occurrence** of this pattern.

A permutation σ **contains** a permutation (or a **pattern**) $\pi = \pi(1), \dots, \pi(k) \in S_k$ if there exist indices $c_1 < \dots < c_k$ such that $\sigma(c_1) \dots \sigma(c_k)$ is order-isomorphic to π . The set of points of indices c_1, \dots, c_k is an **occurrence** of the π .


Pattern (classic)



$\sigma = 413526$ contains the pattern $\pi = 213$.
Positions 1,5,6 is an **occurrence** of this pattern.

A permutation σ **contains** a permutation (or a **pattern**) $\pi = \pi(1), \dots, \pi(k) \in S_k$ if there exist indices $c_1 < \dots < c_k$ such that $\sigma(c_1) \dots \sigma(c_k)$ is order-isomorphic to π . The set of points of indices c_1, \dots, c_k is an **occurrence** of the π .

Pattern (classic)



$\sigma = 413526$ contains the pattern $\pi = 213$.

Positions 1,5,6 is an **occurrence** of this pattern.

$S_n(\pi) :=$ the set of permutations that avoids π .

$S_n(\pi_1, \dots, \pi_k) := \dots$ avoids **all** π_1, \dots, π_k .

A permutation σ **contains** a permutation (or a **pattern**) $\pi = \pi(1), \dots, \pi(k) \in S_k$ if there exist indices $c_1 < \dots < c_k$ such that $\sigma(c_1) \dots \sigma(c_k)$ is order-isomorphic to π . The set of points of indices c_1, \dots, c_k is an **occurrence** of the π .

Pattern avoidance classes

$$S_n(21) = \{Id_n\}$$

Pattern avoidance classes

$$S_n(21) = \{Id_n\}$$

[Knuth 73] : $|S_n(312)| = \frac{1}{n+1} \binom{2n}{n} = 1, 2, 5, 14, 42, 132, \dots$

Pattern avoidance classes

$$S_n(21) = \{Id_n\}$$

[Knuth 73] : $|S_n(312)| = \frac{1}{n+1} \binom{2n}{n} = 1, 2, 5, 14, 42, 132, \dots$

π and τ are **Wilf-equivalent** of $|S_n(\pi)| = |S_n(\tau)|$.

$$|S_n(21)| = |S_n(12)| = 1$$

Pattern avoidance classes


$$S_n(21) = \{Id_n\}$$

[Knuth 73] : $|S_n(312)| = \frac{1}{n+1} \binom{2n}{n} = 1, 2, 5, 14, 42, 132, \dots$

π and τ are **Wilf-equivalent** of $|S_n(\pi)| = |S_n(\tau)|$.

$$|S_n(21)| = |S_n(12)| = 1$$

π and τ are **trivially Wilf-equivalent** if there is a symmetry s of the square such that $\forall \sigma, \sigma \in S(\pi)$ iff $s(\sigma) \in S(s(\tau))$.



Pattern avoidance classes


$$S_n(21) = \{Id_n\}$$

[Knuth 73] : $|S_n(312)| = \frac{1}{n+1} \binom{2n}{n} = 1, 2, 5, 14, 42, 132, \dots$

π and τ are **Wilf-equivalent** of $|S_n(\pi)| = |S_n(\tau)|$.

$$|S_n(21)| = |S_n(12)| = 1$$

π and τ are **trivially Wilf-equivalent** if there is a symmetry s of the square such that $\forall \sigma, \sigma \in S(\pi)$ iff $s(\sigma) \in S(s(\tau))$.



312, 231, 132 and 213 are trivially Wilf-equivalent.

123 and 321 are trivially Wilf-equivalent.

Pattern avoidance classes

TWE: Number of trivially Wilf-Equivalent patterns.

Patterns	<i>TWE</i>	Sequence	Comment
12	2	1, 1, 1, 1, 1, 1, 1, ...	

Pattern avoidance classes

TWE: Number of trivially Wilf-Equivalent patterns.

Patterns	<i>TWE</i>	Sequence	Comment
12	2	1, 1, 1, 1, 1, 1, 1, ...	
12, 21	1	1, 0, 0, 0, 0, 0, 0, ...	

Pattern avoidance classes

TWE: Number of trivially Wilf-Equivalent patterns.

Patterns	<i>TWE</i>	Sequence	Comment
12	2	1, 1, 1, 1, 1, 1, 1, ...	
12, 21	1	1, 0, 0, 0, 0, 0, 0, ...	
312	4	$\frac{1}{n+1} \binom{2n}{n} = 1, 2, 5, 14, 42, 132, \dots$	[Knuth 73]

Pattern avoidance classes

TWE: Number of trivially Wilf-Equivalent patterns.

Patterns	<i>TWE</i>	Sequence	Comment
12	2	1, 1, 1, 1, 1, 1, 1, ...	
12, 21	1	1, 0, 0, 0, 0, 0, 0, ...	
312	4	$\frac{1}{n+1} \binom{2n}{n} = 1, 2, 5, 14, 42, 132, \dots$	[Knuth 73]
123	2		

Pattern avoidance classes

TWE: Number of trivially Wilf-Equivalent patterns.

Patterns	<i>TWE</i>	Sequence	Comment
12	2	1, 1, 1, 1, 1, 1, 1, ...	
12, 21	1	1, 0, 0, 0, 0, 0, 0, ...	
312	4	$\frac{1}{n+1} \binom{2n}{n} = 1, 2, 5, 14, 42, 132, \dots$	[Knuth 73]
123	2	$\frac{1}{n+1} \binom{2n}{n} = 1, 2, 5, 14, 42, 132, \dots$	[Knuth 73]

Pattern avoidance classes

TWE: Number of trivially Wilf-Equivalent patterns.

Patterns	TWE	Sequence	Comment
12	2	1, 1, 1, 1, 1, 1, 1, ...	
12, 21	1	1, 0, 0, 0, 0, 0, 0, ...	
312	4	$\frac{1}{n+1} \binom{2n}{n} = 1, 2, 5, 14, 42, 132, \dots$	[Knuth 73]
123	2	$\frac{1}{n+1} \binom{2n}{n} = 1, 2, 5, 14, 42, 132, \dots$	[Knuth 73]
123, 321	1	1, 2, 4, 4, 0, 0, 0, ...	[Simion 85]
213, 321	4	$1 + \frac{n(n-1)}{2} = 1, 2, 4, 7, 11, 16, 22, \dots$	[Simion 85]
312, 231	2	$2^{n-1} = 1, 2, 4, 8, 16, 32, 64, \dots$	[Simion 85]
231, 132	4	$2^{n-1} = 1, 2, 4, 8, 16, 32, 64, \dots$	[Simion 85]
312, 321	4	$2^{n-1} = 1, 2, 4, 8, 16, 32, 64, \dots$	[Simion 85]

Pattern avoidance classes

TWE: Number of trivially Wilf-Equivalent patterns.


Patterns	TWE	Sequence	Comment
12	2	1, 1, 1, 1, 1, 1, 1, ...	
12, 21	1	1, 0, 0, 0, 0, 0, 0, ...	
312	4	$\frac{1}{n+1} \binom{2n}{n} = 1, 2, 5, 14, 42, 132, \dots$	[Knuth 73]
123	2	$\frac{1}{n+1} \binom{2n}{n} = 1, 2, 5, 14, 42, 132, \dots$	[Knuth 73]
123, 321	1	1, 2, 4, 4, 0, 0, 0, ...	[Simion 85]
213, 321	4	$1 + \frac{n(n-1)}{2} = 1, 2, 4, 7, 11, 16, 22, \dots$	[Simion 85]
312, 231	2	$2^{n-1} = 1, 2, 4, 8, 16, 32, 64, \dots$	[Simion 85]
231, 132	4	$2^{n-1} = 1, 2, 4, 8, 16, 32, 64, \dots$	[Simion 85]
312, 321	4	$2^{n-1} = 1, 2, 4, 8, 16, 32, 64, \dots$	[Simion 85]
213, 132, 123	2	Fibonacci: 1, 2, 3, 5, 8, 13, 21, ...	[Simion 85]
231, 213, 321	8	$n = 1, 2, 3, 4, 5, 6, 7, \dots$	[Simion 85]
312, 132, 213	4	$n = 1, 2, 3, 4, 5, 6, 7, \dots$	[Simion 85]
312, 321, 123	4	1, 2, 3, 1, 0, 0, 0, ...	
321, 213, 123	4	1, 2, 3, 1, 0, 0, 0, ...	
321, 213, 132	2	$n = 1, 2, 3, 4, 5, 6, 7, \dots$	[Simion 85]

patterns and 3-permutations

σ **contains** a pattern π if

patterns and 3-permutations


σ contains a pattern π if there exists a subset of points of σ that is equal (once standardized) to π .



(1432, 3124) contains the pattern (132, 213) .

patterns and 3-permutations

σ **contains** a pattern π if there exists a subset of points of σ that is equal (once standardized) to π .




(1432, 3124) contains the pattern (132, 213) and the pattern 231.

σ **contains** a (2D) pattern π_1 if proj_{xy} , proj_{xz} or proj_{yz} contains π_1 .

patterns and d -permutations

Let $\sigma \in S_n^{d-1}$ -and $\pi \in S_k^{d'-1}$ with $k \leq n$. Then σ **contains the pattern** π , if there exist a **direct** projection $\sigma' = \text{proj}_i(\sigma)$ of dimension d' that contains π .



(1432, 3124) contains the pattern (132, 213) and the pattern 231.

$i := i_1, \dots, i_{d'} \in [d]^{d'}$, the projection proj_i is **direct** if $i_1 < i_2 < \dots < i_{d'}$.
 proj_{xy} is direct. proj_{yx} is **indirect**.

3-pattern avoidance classes

Patterns	TWE	Sequence	Comment
(12, 12)	4	1, 3, 17, 151, 1899, 31711, ...	weak-Bruhat intervals

3-pattern avoidance classes

Patterns	TWE	Sequence	Comment
(12, 12)	4	1, 3, 17, 151, 1899, 31711, ...	weak-Bruhat intervals
(12, 12), (12, 21)	6	$n! = 1, 2, 6, 24, 120 \dots$	$\sigma_1 \Rightarrow \sigma_2$

3-pattern avoidance classes

Patterns	TWE	Sequence	Comment
(12, 12)	4	1, 3, 17, 151, 1899, 31711, ...	weak-Bruhat intervals
(12, 12), (12, 21)	6	$n! = 1, 2, 6, 24, 120, \dots$	$\sigma_1 \Rightarrow \sigma_2$
(12, 12), (12, 21), (21, 12)	4	1, 1, 1, 1, 1, 1, ...	1 diagonal
(12, 12), (12, 21), (21, 12), (21, 21)	1	1, 0, 0, 0, 0, 0, ...	

3-pattern avoidance classes

Patterns	TWE	Sequence	Comment
(12, 12)	4	1, 3, 17, 151, 1899, 31711, ...	weak-Bruhat intervals
(12, 12), (12, 21)	6	$n! = 1, 2, 6, 24, 120, \dots$	$\sigma_1 \Rightarrow \sigma_2$
(12, 12), (12, 21), (21, 12)	4	1, 1, 1, 1, 1, 1, ...	1 diagonal
(12, 12), (12, 21), (21, 12), (21, 21)	1	1, 0, 0, 0, 0, 0, ...	
(123, 123)	4	1, 4, 35, 524, 11774, 366352, ...	<i>new</i>
(123, 132)	24	1, 4, 35, 524, 11768, 365558, ...	<i>new</i>
(132, 213)	8	1, 4, 35, 524, 11759, 364372, ...	<i>new</i>

3-pattern avoidance classes

Patterns	TWE	Sequence	Comment
(12, 12)	4	1, 3, 17, 151, 1899, 31711, ...	weak-Bruhat intervals
(12, 12), (12, 21)	6	$n! = 1, 2, 6, 24, 120, \dots$	$\sigma_1 \Rightarrow \sigma_2$
(12, 12), (12, 21), (21, 12)	4	1, 1, 1, 1, 1, 1, ...	1 diagonal
(12, 12), (12, 21), (21, 12), (21, 21)	1	1, 0, 0, 0, 0, 0, ...	
(123, 123)	4	1, 4, 35, 524, 11774, 366352, ...	<i>new</i>
(123, 132)	24	1, 4, 35, 524, 11768, 365558, ...	<i>new</i>
(132, 213)	8	1, 4, 35, 524, 11759, 364372, ...	<i>new</i>
(12, 12), (132, 312)	48	$(n+1)^{n-1} = 1, 3, 16, 125, 1296, \dots$	[Atkinson et al. 93,95]
(12, 12), (123, 321)	12	1, 3, 16, 124, 1262, 15898, ...	distributive lattices inter.
(12, 12), (231, 312)	8	1, 3, 16, 122, 1188, 13844, ...	A295928?

3-pattern avoidance classes

[A295928](#) Number of triangular matrices $T(n,i,k)$, $k \leq i \leq n$, with entries "0" or "1" +40¹ with the property that each triple $\{T(n,i,k), T(n,i,k+1), T(n,i-1,k)\}$ containing a single "0" can be successively replaced by $\{1, 1, 1\}$ until finally no "0" entry remains.

1, 3, 16, 122, 1188, 13844, 185448, 2781348, 45868268 ([list](#); [graph](#); [refs](#); [listen](#); [history](#); [text](#); [internal format](#))

OFFSET 1,2

COMMENTS A triple $\{T(n,i,k), T(n,i,k+1), T(n,i-1,k)\}$ will be called a primitive triangle. It is easy to see that $b(n) = n(n-1)/2$ is the number of such triangles. At each step, exactly one primitive triangle is completed (replaced by $\{1, 1, 1\}$). So there are $b(n)$ "0"- and n "1"-terms. Thus the starting matrix has no complete primitive triangle. Furthermore, any triangular submatrix $T(m,i,k)$, $k \leq i \leq m < n$ cannot have more than m "1"-terms because otherwise it would have less "0"-terms than primitive triangles. The replacement of at least one "0"-term would complete more than one primitive triangle. This has been excluded.
So $T(n, i, k)$ is a special case of $U(n, i, k)$, described in [A101481](#): $a(n) < \text{A101481}(n+1)$.

A start matrix may serve as a pattern for a number wall used on worksheets for elementary mathematics, see link "Number walls". That is why I prefer the more descriptive name "fill matrix".

The algorithm for the sequence is rather slow because each start matrix is constructed separately. There exists a faster recursive algorithm which produces the same terms and therefore is likely to be correct, but it is based on a conjecture. For the theory of the recurrence, see "Recursive aspects of fill matrices". Probable extension $a(10)-a(14)$: 821096828, 15804092592, 324709899276, 7081361097108, 163179784397820.

The number of fill matrices with n rows and all "1"- terms concentrated on the last two rows, is [A001960](#)(n).

See link "Reconstruction of a sequence".

LINKS

[Table of \$n, a\(n\)\$ for \$n=1..9\$](#) .

Gerhard Kirchner, [Recursive aspects of fill matrices](#)

Gerhard Kirchner, [Number walls](#)

Gerhard Kirchner, [VB-program](#)

Gerhard Kirchner, [Reconstruction of a sequence](#)

Ville Salo, [Cutting Corners](#), arXiv:2002.08730 [math.DS], 2020.

2-pattern avoidance classes

Patterns	TWE	Sequence	Comment
12	1	1, 0, 0, 0, 0, ⋯	unvoidable pattern

2-pattern avoidance classes

Patterns	TWE	Sequence	Comment
12	1	1, 0, 0, 0, 0, ⋯	unvoidable pattern
21	1	1, 1, 1, 1, 1, ⋯	1 diagonal

2-pattern avoidance classes

Patterns	TWE	Sequence	Comment
12	1	1, 0, 0, 0, 0, ...	unvoidable pattern
21	1	1, 1, 1, 1, 1, ...	1 diagonal
123	1	1, 4, 20, 100, 410, 1224, 2232, ...	<i>new</i>
132	2	1, 4, 21, 116, 646, 3596, 19981, ...	<i>new</i>
231	2	1, 4, 21, 123, 767, 4994, 33584, ...	<i>new</i>
321	1	1, 4, 21, 128, 850, 5956, 43235, ...	<i>new</i>

2-pattern avoidance classes

Patterns	TWE	Sequence	Comment
12	1	1, 0, 0, 0, 0, ...	unvoidable pattern
21	1	1, 1, 1, 1, 1, ...	1 diagonal
123	1	1, 4, 20, 100, 410, 1224, 2232, ...	<i>new</i>
132	2	1, 4, 21, 116, 646, 3596, 19981, ...	<i>new</i>
231	2	1, 4, 21, 123, 767, 4994, 33584, ...	<i>new</i>
321	1	1, 4, 21, 128, 850, 5956, 43235, ...	<i>new</i>
123, 132	2	1, 4, 8, 8, 0, 0, 0, ...	
123, 231	2	1, 4, 9, 6, 0, 0, 0, ...	
123, 321	1	1, 4, 8, 0, 0, 0, 0, ...	
132, 213	1	1, 4, 12, 28, 58, 114, 220, ...	[Sun22+]
132, 231	4	1, 4, 12, 32, 80, 192, 448, ...	A001787 [Sun22+]
132, 321	2	1, 4, 12, 27, 51, 86, 134, ...	A047732 [Sun22+]
231, 312	1	1, 4, 10, 28, 76, 208, 568, ...	A026150 [Sun22+]
231, 321	2	1, 4, 12, 36, 108, 324, 972, ...	A003946 [Sun22+]

2-pattern avoidance classes

A001787 $a(n) = n^*2^{(n-1)}$.
(Formerly M3444 N1398)

407

0, 1, 4, 12, 32, 80, 192, 448, 1024, 2304, 5120, 11264, 24576, 53248, 114688, 245760, 524288, 1114112, 2359296, 4980736, 10485760, 22020096, 46137344, 96468992, 201326592, 419430400, 872415232, 1811939328, 3758096384, 7784628224, 16106127360, 33285996544 ([list](#); [graph](#); [refs](#); [listen](#);

A047732 First differences are [A005563](#).

1

1, 4, 12, 27, 51, 86, 134, 197, 277, 376, 496, 639, 807, 1002, 1226, 1481, 1769, 2092, 2452, 2851, 3291, 3774, 4302, 4877, 5501, 6176, 6904, 7687, 8527, 9426, 10386, 11409, 12497, 13652, 14876, 16171, 17539, 18982, 20502, 22101, 23781, 25544, 27392, 29327 ([list](#); [graph](#); [refs](#); [listen](#); [history](#); [text](#); [internal format](#))

OFFSET 0,2

COMMENTS Number of 3-permutations of n elements avoiding the patterns 132, 321. See Bonichon and Sun. - [Michel Marcus](#), Aug 19 2022

LINKS Vincenzo Librandi, [Table of n, a\(n\) for n = 0..1000](#)
Nicolas Bonichon and Pierre-Jean Morel, [Baxter d-permutations and other pattern avoiding classes](#), arXiv:2202.12677 [math.CO], 2022.

Nathan Sun, [On d-permutations and Pattern Avoidance Classes](#), arXiv:2208.08506 [math.CO], 2022.

[Index entries for linear recurrences with constant coefficients](#), signature (4,-6,4, -1).

FORMULA $a(n) = A051925(n+1) + 1$. - [Alex Ratshnyak](#), Jun 27 2012

From [Vincenzo Librandi](#), Jun 28 2012: (Start)

G.f.: $(1 + 2*x^2 - x^3)/(1-x)^4$.

$a(n) = 4*a(n-1) - 6*a(n-2) + 4*a(n-3) - a(n-4)$.

A026150 $a(0) = a(1) = 1$; $a(n+2) = 2^*a(n+1) + 2^*a(n)$.

53

1, 1, 4, 10, 28, 76, 208, 568, 1552, 4240, 11584, 31648, 86464, 236224, 645376, 1763200, 4817152, 13160704, 35955712, 98232832, 268377088, 733219840, 2003193856, 5472827392, 14952042496, 40849739776 ([list](#); [graph](#); [refs](#); [listen](#); [history](#); [text](#); [internal format](#))

A003946 Expansion of $(1+x)/(1-3*x)$.

127

1, 4, 12, 36, 108, 324, 972, 2916, 8748, 26244, 78732, 236196, 708588, 2125764, 6377292, 19131876, 57395628, 172186884, 516560652, 1549681956, 4649045868, 13947137604, 41841412812, 125524238436, 376572715308, 1129718145924 ([list](#); [graph](#); [refs](#); [listen](#); [history](#); [text](#); [internal format](#))

2– and 3-patterns avoidance classes

Patterns	TWE	Sequence	Comment
12, ($\textcolor{blue}{12}$, $\textcolor{red}{12}$)	1	1, 0, 0, 0, 0, \cdots	12
12, ($\textcolor{blue}{21}$, $\textcolor{red}{12}$)	3	1, 0, 0, 0, 0, \cdots	12
21, ($\textcolor{blue}{12}$, $\textcolor{red}{12}$)	1	1, 0, 0, 0, 0, \cdots	
21, ($\textcolor{blue}{21}$, $\textcolor{red}{12}$)	3	1, 1, 1, 1, 1, \cdots	21
123, ($\textcolor{blue}{12}$, $\textcolor{red}{12}$)	1	1, 3, 14, 70, 288, 822, 1260, \cdots	<i>new</i>
123, ($\textcolor{blue}{12}$, $\textcolor{red}{21}$)	3	1, 3, 6, 6, 0, 0, 0, \cdots	
132, ($\textcolor{blue}{12}$, $\textcolor{red}{12}$)	2	1, 3, 11, 41, 153, 573, 2157, \cdots	A281593?
132, ($\textcolor{blue}{12}$, $\textcolor{red}{21}$)	6	1, 3, 11, 43, 173, 707, 2917, \cdots	A026671?
231, ($\textcolor{blue}{12}$, $\textcolor{red}{12}$)	2	1, 3, 9, 26, 72, 192, 496, \cdots	A072863?
231, ($\textcolor{blue}{12}$, $\textcolor{red}{21}$)	4	1, 3, 11, 44, 186, 818, 3706, \cdots	<i>new</i>
231, ($\textcolor{blue}{21}$, $\textcolor{red}{12}$)	2	1, 3, 12, 55, 273, 1428, 7752, \cdots	A001764?
321, ($\textcolor{blue}{12}$, $\textcolor{red}{12}$)	1	1, 3, 2, 0, 0, 0, 0, \cdots	
321, ($\textcolor{blue}{12}$, $\textcolor{red}{21}$)	3	1, 3, 11, 47, 221, 1113, 5903, \cdots	A217216?

2– and 3-patterns avoidance classes

A281593	$a(n) = b(n) - \sum_{j=0..n-1} b(n)$ with $b(n) = \text{binomial}(2*n, n)$.	1
	1, 1, 3, 11, 41, 153, 573, 2157, 8163, 31043, 118559, 454479, 1747771, 6740059, 26055459, 100939779, 391785129, 1523230569, 5931153429, 23126146629, 90282147849, 352846964649, 1380430179489, 5405662979649, 21186405207549, 83101804279101, 326199124351701 (list ; graph ; refs);	
A026671	Number of lattice paths from (0,0) to (n,n) with steps (0,1), (1,0) and, when on the diagonal, (1,1).	21
	1, 3, 11, 43, 173, 707, 2917, 12111, 50503, 211263, 885831, 3720995, 15652239, 65913927, 277822147, 1171853635, 4945846997, 20884526283, 88224662549, 372827899079, 1576001732485, 6663706588179, 28181895551161, 119208323665543, 584329070986033, 2133944799315027 (list ; graph ; refs);	
A072863	$a(n) = 2^{(n-3)*(n^2+2*3*n+8)}$.	7
	1, 3, 9, 26, 72, 192, 496, 1248, 3072, 7424, 17664, 41472, 96256, 221184, 503808, 1138688, 2555904, 5701632, 12648448, 27918336, 61341696, 134217728, 292552704, 635437056, 1375731712, 2969567232, 6392119296, 13723762688 (list ; graph ; refs ; listen ; history ; text ; internal format)	
A001764	$a(n) = \text{binomial}(3*n, n)/(2*n+1)$ (enumerates ternary trees and also noncrossing trees). (Formerly M2926 N1174)	448
	1, 1, 3, 12, 55, 273, 1428, 7752, 43263, 246675, 1430715, 8414640, 50067108, 300830572, 1822766520, 11124755664, 68328754959, 422030545335, 2619631042665, 16332922290300, 102240109897695, 642312451217745, 4048514844039120, 25594403741131680, 162250238001816900 (list ;	
A217216	Dimension of algebraic generators of the algebra "Baxter" of order n.	0
	0, 1, 1, 3, 11, 47, 221, 1113, 5903, 32607, 186143, 1092015, 6555515, 40137219, 249984481, 1588468321, 10125395007, 65639436955, 430048061915, 2844592155631, 18979693010495, 127641472658231, 864645413540671, 5896221199266519, 40455246946190079 (list ; graph ; refs ; listen ;	

Separable permutations Sep_n

direct sum $\sigma \oplus \pi$: add π in the top right corner of σ .

A Young diagram consisting of two parts. On the left, a green rectangular box contains three dots at positions (1,1), (1,2), and (2,1). On the right, a purple L-shaped hook shape is attached to the top-right corner of the green box, containing two dots at (2,2) and (3,1). The total shape has 5 columns and 4 rows, with labels above it: 1, 3, 2, 5, 4.

$$132 \oplus 21 = 13254$$

skew sum $\sigma \ominus \pi$: add π in the bottom right corner of σ .


A Young diagram consisting of two parts. On the left, a green rectangular box contains three dots at positions (1,1), (1,2), and (2,1). On the right, a purple L-shaped hook shape is attached to the bottom-right corner of the green box, containing two dots at (2,2) and (3,1). The total shape has 5 columns and 4 rows, with labels above it: 3, 5, 4, 2, 1.

$$132 \ominus 21 = 35421$$

Separable permutations Sep_n


direct sum $\sigma \oplus \pi$: add π in the top right corner of σ .

$$132 \oplus 21 = 13254$$




skew sum $\sigma \ominus \pi$: add π in the bottom right corner of σ .

$$132 \ominus 21 = 35421$$



separable: size 1 or a direct/skew sum separable permutations.



$$643512 = 1 \ominus ((1 \ominus 1) \oplus 1) \ominus (1 \oplus 1).$$

Separable permutations Sep_n

direct sum $\sigma \oplus \pi$: add π in the top right corner of σ .

$$132 \oplus 21 = 13254$$

skew sum $\sigma \ominus \pi$: add π in the bottom right corner of σ .

$$132 \ominus 21 = 35421$$

separable: size 1 or a direct/skew sum separable permutations.

Separable permutations Sep_n

separable: size 1 or a direct/skew sum separable permutations.

[Brightwell 92]:


$$|Sep_n| = \frac{1}{n-1} \sum_{k=0}^{n-2} \binom{n-1}{k} \binom{n-1}{k+1} 2^{n-k-1}.$$

Separable permutations Sep_n

separable: size 1 or a direct/skew sum separable permutations.
[Brightwell 92]:

$$|Sep_n| = \frac{1}{n-1} \sum_{k=0}^{n-2} \binom{n-1}{k} \binom{n-1}{k+1} 2^{n-k-1}.$$

$Sep_n = S_n(2413, 3142)$ [Bose Buss Lubiw 98]




Separable d -permutations [Atkinson Mansour 10]

d -sum:


$$\pi \oplus^{(+++)} \sigma$$

$$[[1, 3, 2, 5, 4], [1, 3, 2, 4, 5]]$$




$$\pi \oplus^{(++-)} \sigma$$

$$[[1, 3, 2, 5, 4], [3, 5, 4, 1, 2]]$$




$$\pi \oplus^{(+--)} \sigma$$

$$[[1, 5, 4, 2, 1], [1, 3, 2, 4, 5]]$$



$$\pi \oplus^{(--)} \sigma$$

$$[[3, 5, 4, 2, 1], [3, 5, 4, 1, 2]]$$



Let σ and π two d -permutations and $\text{dir} \in \{+, -\}^d$. The **d -sum** with respect to direction dir is :

$$\sigma \oplus^{\text{dir}} \pi := \bar{\sigma}_2 \oplus_2^{\text{dir}} \bar{\pi}_2, \dots, \bar{\sigma}_d \oplus_d^{\text{dir}} \bar{\pi}_d,$$


where \oplus_i^{dir} is \oplus if $\text{dir}_i = +$ and \ominus if $\text{dir}_i = -$.

Separable d -permutations [Atkinson Mansour 10]

d -sum:


$$\pi \oplus^{(+++)} \sigma$$

$[[1, 3, 2, 5, 4], [1, 3, 2, 4, 5]]$




$$\pi \oplus^{(+-+)} \sigma$$

$[[1, 3, 2, 5, 4], [3, 5, 4, 1, 2]]$




$$\pi \oplus^{(+--)} \sigma$$

$[[3, 5, 4, 2, 1], [1, 3, 2, 4, 5]]$



$$\pi \oplus^{(--)} \sigma$$



$[[3, 5, 4, 2, 1], [3, 5, 4, 1, 2]]$




separable d -permutation: size 1 or a d -sum separable permutations.

$$p_1 = (132, 132) = (1, 1) \oplus^{(+++)} ((1, 1) \oplus^{(+-+)} (1, 1))$$

Separable d -permutations [Atkinson Mansour 10]





Separable d -permutations [Atkinson Mansour 10]



[Atkinson Mansour 10]

$$Sep_n^{d-1} = S_n^{d-1}(2413, 3142, \text{Sym}((\textcolor{blue}{132}, \textcolor{red}{213})))$$

Separable d -permutations [Atkinson Mansour 10]



[Atkinson Mansour 10]

$$Sep_n^{d-1} = S_n^{d-1}(2413, 3142, \text{Sym}((\textcolor{blue}{132}, \textcolor{red}{213})))$$

[Atkinson Mansour 10]

$$|Sep_n^{d-1}| = \frac{1}{n-1} \sum_{k=0}^{n-2} \binom{n-1}{k} \binom{n-1}{k+1} (2^{d-1} - 1)^k (2^{d-1})^{n-k-1}.$$

random generation and limit

THE BROWNIAN LIMIT OF SEPARABLE PERMUTATIONS

FRÉDÉRIQUE BASSINO, MATHILDE BOUVEL, VALENTIN FÉRAY, LUCAS GERIN, AND ADELINE PIERROT

ABSTRACT. We study uniform random permutations in an important class of pattern-avoiding permutations: the separable permutations. We describe the asymptotics of the number of occurrences of any fixed given pattern in such a random permutation in terms of the Brownian excursion. In the recent terminology of permutons, our work can be interpreted as the convergence of uniform random separable permutations towards a “Brownian separable permuton”.




FIGURE 1. Two uniform random separable permutations of sizes respectively $n = 204523$ and $n = 903\,073$ (a permutation σ is represented here with its *diagram*: for every $i \leq n$, there is a dot at coordinates (i, σ_i)).

random generation and limit

THE BROWNIAN LIMIT OF SEPARABLE PERMUTATIONS

FRÉDÉRIQUE BASSINO, MATHILDE BOUVEL, VALENTIN FÉRAY, LUCAS GERIN, AND ADELINE PIERROT

ABSTRACT. We study uniform random permutations in an important class of pattern-avoiding permutations: the separable permutations. We describe the asymptotics of the number of occurrences of any fixed given pattern in such a random permutation in terms of the Brownian excursion. In the recent terminology of permutons, our work can be interpreted as the convergence of uniform random separable permutations towards a “Brownian separable permuton”.





FIGURE 1. Two uniform random separable permutations of sizes respectively $n = 204\,523$ and $n = 903\,073$ (a permutation σ is represented here with its *diagram*: for every $i \leq n$, there is a dot at coordinates (i, σ_i)).

Random separable 3-permutations of size ≈ 100000 .(with UsainBoltz)

Vincular Patterns and Baxter permutations



vincular pattern: a pattern where some entries must be consecutive in the permutation (**adjacencies**). Ex: $2413|_2$ and $3142|_2$



A vincular pattern $\pi|_X$, $\pi \in S_k$, $X \subseteq [k - 1]$. $\sigma \in S_n$ contains $\pi|_X$, $\sigma_{i_1}, \sigma_{i_2} \dots \sigma_{i_k} \equiv \pi$ and that $i_{j+1} = i_j + 1 \forall j \in X$.

Vincular Patterns and Baxter permutations



vincular pattern: a pattern where some entries must be consecutive in the permutation (**adjacencies**). Ex: $2413|_2$ and $3142|_2$



A vincular pattern $\pi|_X$, $\pi \in S_k$, $X \subseteq [k - 1]$. $\sigma \in S_n$ contains $\pi|_X$, $\sigma_{i_1}, \sigma_{i_2} \dots \sigma_{i_k} \equiv \pi$ and that $i_{j+1} = i_j + 1 \forall j \in X$.

Vincular Patterns and Baxter permutations

vincular pattern: a pattern where some entries must be consecutive in the permutation (**adjacencies**). Ex: $2413|_2$ and $3142|_2$




Baxter permutations: $B_n := S_n(2413|_2, 3142|_2)$.

A vincular pattern $\pi|_X$, $\pi \in S_k$, $X \subseteq [k - 1]$. $\sigma \in S_n$ contains $\pi|_X$, $\sigma_{i_1}, \sigma_{i_2} \dots \sigma_{i_k} \equiv \pi$ and that $i_{j+1} = i_j + 1 \forall j \in X$.

Generalized vincular Patterns


generalized vincular pattern : horizontal and/or vertical adjacencies. Ex: $2413|_{2,2}$ and $3142|_{2,2}$.



$$B_n = S_n(2413|_{2,2}, 3142|_{2,2}).$$

Generalized vincular Patterns

generalized vincular pattern : horizontal and/or vertical adjacencies. Ex: $2413|_{2,2}$ and $3142|_{2,2}$.




$$B_n = S_n(2413|_{2,2}, 3142|_{2,2}).$$

Natural extension of generalized vincular patterns to
 d -permutations...

Generalized vincular Patterns

generalized vincular pattern : horizontal and/or vertical adjacencies. Ex: $2413|_{2,2}$ and $3142|_{2,2}$.




$$B_n = S_n(2413|_{2,2}, 3142|_{2,2}).$$

Natural extension of generalized vincular patterns to
 d -permutations...

But what could be a Baxter d -permutation...


Well sliced permutations

A **slice** is a rectangle defined by two adjacent points. **type**: horizontal or vertical. The **direction** of a slice is: **+** or **-**.





Well sliced permutations

A **slice** is a rectangle defined by two adjacent points. **type**: horizontal or vertical. The **direction** of a slice is: **+** or **-**.





Well sliced permutations

A **slice** is a rectangle defined by two adjacent points. **type**: horizontal or vertical. The **direction** of a slice is: **+** or **-**.



Well sliced permutations


A **slice** is a rectangle defined by two adjacent points. **type**: horizontal or vertical. The **direction** of a slice is: **+** or **-**.




well-sliced: each vertical slice intersects exactly 1 horizontal slice and 2 intersecting slices share the same direction.

Well sliced permutations

A **slice** is a rectangle defined by two adjacent points. **type**: horizontal or vertical. The **direction** of a slice is: **+** or **-**.



not well sliced





well sliced

well-sliced: each vertical slice intersects exactly 1 horizontal slice and 2 intersecting slices share the same direction.


Well sliced permutations

A **slice** is a rectangle defined by two adjacent points. **type**: horizontal or vertical. The **direction** of a slice is: **+** or **-**.







well-sliced: each vertical slice intersects exactly 1 horizontal slice and 2 intersecting slices share the same direction.

Proposition: Baxter \Leftrightarrow well-sliced







Well-sliced d -permutations

The **direction** of a slice is: ++, +- , -+ , -. The **type** of a slice is: x, y or z.



Well-sliced d -permutations


The **direction** of a slice is: $++$, $+ -$, $- +$, $- -$. The **type** of a slice is: x , y or z .



well-sliced: each slice intersects exactly 1 slice of each type and 2 intersecting slices share the same direction.

Well-sliced d -permutations





The **direction** of a slice is: ++, +- , -+, -. The **type** of a slice is: x, y or z.




well-sliced: each slice intersects exactly 1 slice of each type and 2 intersecting slices share the same direction.

Baxter d -permutation

A **Baxter d -permutation** is a d -permutation such that each of its $d' \leq d$ projection is well-sliced.



Well-sliced d -permutations vs Baxter d -permutation



well-sliced but not Baxter.

Baxter d -permutation characterisation

Theorem

$$B_n^{d-1} = S_n^{d-1}(\text{Sym}(2413|_{2,2}), \text{Sym}((\textcolor{blue}{3}12, \textcolor{red}{2}13)|_{1,\textcolor{red}{2},\dots}), \\ \text{Sym}((\textcolor{blue}{3}412, \textcolor{red}{1}432)|_{2,\textcolor{red}{2},\dots}), \text{Sym}((\textcolor{blue}{2}143, \textcolor{red}{1}423)|_{2,\textcolor{red}{2},\dots})).$$

Baxter d -permutation enumration


$$|B_n^{d-1}|$$

$n \setminus d$	2	3	4	5
1	1	1	1	1
2	2	4	8	16
3	6	28	120	496
4	22	260	2440	20816
5	92	2872	59312	1035616
6	422	35620		
7	2074	479508		

open problem: Enumeration formula?

Baxter / Complete Baxter / Anti-Baxter permutations

dots: intersection of slices





squares: Baxter permutations

squares + dots : Complete Baxter permutations

dots: Anti-Baxter permutation

Baxter / Complete Baxter / Anti-Baxter permutations

dots: intersection of slices



[B. Bousquet Fusy 10] Baxter permutations \leftrightarrow plane bipolar orientations.

Baxter / Complete Baxter / Anti-Baxter
 d -permutations

Anti-Baxter d -permutation enumeration

$$|A_n^{d-1}|$$

$n \setminus d$	2	3	4	5
1	1	1	1	1
2	2	4	8	16
3	6	36	216	1296
4	22	444	7096	
5	88	5344		
6	374	64460		
7	1668			

open problem: Enumeration formula?

conclusion/perspectives

- general framework for pattern avoidance in d -permutations
+ easy results
- generalisation of Baxter permutation
- work in progress (with Thomas Muller and Adrian Tanasa): links between 2^{d-1} floorplans and d -permutations.
- lot of open problems
- implementation available

plmlab.math.cnrs.fr/bonichon/multipermutation

conclusion/perspectives

- general framework for pattern avoidance in d -permutations
+ easy results
- generalisation of Baxter permutation
- work in progress (with Thomas Muller and Adrian Tanasa): links between 2^{d-1} floorplans and d -permutations.
- lot of open problems
- implementation available

`plmlab.math.cnrs.fr/bonichon/multipermutation`

Have fun !