

An unexpected symmetry on graphs

Florian Fürnsinn, Moritz Gangl and Martin Rubey

Universität & Technische Universität Wien

Bordeaux 2025





[Navigation](#)

[All Collections](#)

[All Maps](#)

[All Statistics](#)

[Contribute Map](#)

[Contribute Statistic](#)

[Usage](#)

[Contributors](#)

[Citations](#)

[Interface](#)

— [database and search engine for combinatorial statistics and maps](#) —

go to **statistic finder**

[start](#)

go to **map finder**

[start](#)

search for **keyword**

[start](#)

[sign up for irregular FindStat news](#)
[contact the FindStat team at info@findstat.org](mailto:info@findstat.org)

Let \mathcal{C}_n be a finite collection of objects, e.g.

- ▶ permutations in \mathfrak{S}_n ,
- ▶ Dyck paths of semilength n ,
- ▶ (simple, unlabelled) graphs with n vertices, . . .

A *statistic* is a function $s : \mathcal{C}_n \rightarrow \mathbb{Z}$, its *distribution* is $\sum_{c \in \mathcal{C}_n} q^{s(c)}$.

Let \mathcal{C}_n be a finite collection of objects, e.g.

- ▶ permutations in \mathfrak{S}_n ,
- ▶ Dyck paths of semilength n ,
- ▶ (simple, unlabelled) graphs with n vertices, . . .

A *statistic* is a function $s : \mathcal{C}_n \rightarrow \mathbb{Z}$, its *distribution* is $\sum_{c \in \mathcal{C}_n} q^{s(c)}$.

In www.findstat.org, we have

Permutations	~ 400 statistics	18 bijections
--------------	------------------	---------------

Dyck paths	~ 300 statistics	24 bijections
------------	------------------	---------------

...

Let \mathcal{C}_n be a finite collection of objects, e.g.

- ▶ permutations in \mathfrak{S}_n ,
- ▶ Dyck paths of semilength n ,
- ▶ (simple, unlabelled) graphs with n vertices, . . .

A *statistic* is a function $s : \mathcal{C}_n \rightarrow \mathbb{Z}$, its *distribution* is $\sum_{c \in \mathcal{C}_n} q^{s(c)}$.

In www.findstat.org, we have

Permutations	~ 400 statistics	18 bijections
Dyck paths	~ 300 statistics	24 bijections
...		
(simple, unlabelled) graphs	~ 300 statistics	2 bijections

A *statistic* is a function $s : \mathcal{C}_n \rightarrow \mathbb{Z}$, its *distribution* is $\sum_{c \in \mathcal{C}_n} q^{s(c)}$.
If s_1 and s_2 are equidistributed, $s_1 \sim s_2$, there must be a bijection

$$\begin{aligned}\phi : \mathcal{C}_n &\rightarrow \mathcal{C}_n \\ s_1(\phi(e)) &= s_2(e).\end{aligned}$$

A *statistic* is a function $s : \mathcal{C}_n \rightarrow \mathbb{Z}$, its *distribution* is $\sum_{c \in \mathcal{C}_n} q^{s(c)}$.
If s_1 and s_2 are equidistributed, $s_1 \sim s_2$, there must be a bijection

$$\begin{aligned}\phi : \mathcal{C}_n &\rightarrow \mathcal{C}_n \\ s_1(\phi(e)) &= s_2(e).\end{aligned}$$

Many statistics on Dyck paths or permutations have the same distribution, e.g.

Permutations $maj \sim inv$
Dyck paths $height \sim bounce\ count$

No such (non-trivial) equidistributions were known for graphs.

A *statistic* is a function $s : \mathcal{C}_n \rightarrow \mathbb{Z}$, its *distribution* is $\sum_{c \in \mathcal{C}_n} q^{s(c)}$.
If s_1 and s_2 are equidistributed, $s_1 \sim s_2$, there must be a bijection

$$\begin{aligned}\phi : \mathcal{C}_n &\rightarrow \mathcal{C}_n \\ s_1(\phi(e)) &= s_2(e).\end{aligned}$$

Many statistics on Dyck paths or permutations have the same distribution, e.g.

Permutations *maj* \sim *inv*
Dyck paths *height* \sim *bounce count*

No such (non-trivial) equidistributions were known for graphs.
However, **FindStat** found

#graphs without leaves = #graphs with distinct neighbourhoods

mating

Theorem (Kilibarda 2007, Gessel & Li 2011).

#graphs without leaves = #graphs with distinct neighbourhoods

mating

Theorem (Kilibarda 2007, Gessel & Li 2011).

#graphs without leaves = #graphs with distinct neighbourhoods

Theorem (Read 1989). The generating function for graphs with distinct neighbourhoods is

$$\sum_{\lambda} \frac{2^{X_{\lambda}}}{z_{\lambda}} x^{|\lambda|} \prod_i (1 - x^{\lambda_i}),$$

where $z_{\lambda} = 1^{m_1} m_1! 2^{m_2} m_2! \dots$ for $\lambda = (1^{m_1}, 2^{m_2}, \dots)$ and

$$X_{\lambda} = \sum_{i < j} \gcd(\lambda_i, \lambda_j) + \sum_i \left\lfloor \frac{\lambda_i}{2} \right\rfloor.$$

mating

Definition (Sumner 1973, Bull & Pease 1989).

The *neighbourhood of u* is $N(u) = \{v \mid (u, v) \in E(G)\}$,
the *closed neighbourhood of u* is $\bar{N}(u) = \{u\} \cup N(u)$.

mating

Definition (Sumner 1973, Bull & Pease 1989).

The *neighbourhood of u* is $N(u) = \{v \mid (u, v) \in E(G)\}$,

the *closed neighbourhood of u* is $\overline{N}(u) = \{u\} \cup N(u)$.

A graph is *point determining* or *mating*, if $N(u) \neq N(v)$,

it is *point distinguishing* or *co-mating*, if $\overline{N}(u) \neq \overline{N}(v)$,

for any pair of vertices $u \neq v$.


Theorem (Kilibarda 2007, Gessel & Li 2011).

$\#$ graphs without leaves = $\#$ mating graphs = $\#$ co-mating graphs


tufts and siblings

Definition (Alexandersson & R. 2022). The *tuft number* $t(G)$ is the maximal number of leaves adjacent to the same vertex.


a *tuft*:




$t(G) =$




$t(G) =$




$t(G) =$



$t(G) =$



$t(G) =$




$t(G) =$


tufts and siblings

Definition (Alexandersson & R. 2022). The *tuft number* $t(G)$ is the maximal number of leaves adjacent to the same vertex.


a *tuft*:




$$t(G) = 3$$




$$t(G) = 1$$




$$t(G) = 0$$



$$t(G) = 1$$



$$t(G) = 0$$




$$t(G) = 0$$

tufts and siblings


Definition (Alexandersson & R. 2022). The *tuft number* $t(G)$ is the maximal number of leaves adjacent to the same vertex.

The *sibling number* is the maximal number of siblings of a vertex:


$$s(G) = \max_{v \in V(G)} |\{u \in V(G) \mid \bar{N}(u) = \bar{N}(v)\}| - 1.$$




$$s(G) = \\ t(G) = 3$$




$$s(G) = \\ t(G) = 1$$




$$s(G) = \\ t(G) = 0$$



$$s(G) = \\ t(G) = 1$$



$$s(G) = \\ t(G) = 0$$




$$s(G) = \\ t(G) = 0$$

tufts and siblings


Definition (Alexandersson & R. 2022). The *tuft number* $t(G)$ is the maximal number of leaves adjacent to the same vertex.

The *sibling number* is the maximal number of siblings of a vertex:


$$s(G) = \max_{v \in V(G)} |\{u \in V(G) \mid \bar{N}(u) = \bar{N}(v)\}| - 1.$$




$$s(G) = 0 \\ t(G) = 3$$




$$s(G) = 0 \\ t(G) = 1$$




$$s(G) = 0 \\ t(G) = 0$$



$$s(G) = 1 \\ t(G) = 1$$



$$s(G) = 1 \\ t(G) = 0$$



$$s(G) = 3 \\ t(G) = 0$$

tufts and siblings


Definition (Alexandersson & R. 2022). The *tuft number* $t(G)$ is the maximal number of leaves adjacent to the same vertex.

The *sibling number* is the maximal number of siblings of a vertex:


$$s(G) = \max_{v \in V(G)} |\{u \in V(G) \mid \bar{N}(u) = \bar{N}(v)\}| - 1.$$

Conjecture (Alexandersson & R. 2022).


$$(s, t) \sim (t, s)$$




$$s(G) = 0 \\ t(G) = 3$$




$$s(G) = 0 \\ t(G) = 1$$




$$s(G) = 0 \\ t(G) = 0$$



$$s(G) = 1 \\ t(G) = 1$$



$$s(G) = 1 \\ t(G) = 0$$



$$s(G) = 3 \\ t(G) = 0$$

tufts and siblings


Definition (Alexandersson & R. 2022). The *tuft number* $t(G)$ is the maximal number of leaves adjacent to the same vertex.

The *sibling number* is the maximal number of siblings of a vertex:

$$s(G) = \max_{v \in V(G)} |\{u \in V(G) \mid \bar{N}(u) = \bar{N}(v)\}| - 1.$$


Theorem (Fürnsinn & Gangl & R. 2024).

$$(s, t) \sim (t, s)$$




$$s(G) = 0$$

$$t(G) = 3$$




$$s(G) = 0$$

$$t(G) = 1$$




$$s(G) = 0$$

$$t(G) = 0$$




$$s(G) = 1$$

$$t(G) = 1$$



$$s(G) = 1$$

$$t(G) = 0$$




$$s(G) = 3$$

$$t(G) = 0$$

first idea: look at small examples and find a bijection

Fact. In graphs other than K_2 , siblings cannot be adjacent to leaves.

Contract tufts with n leaves to a vertex T_n
and groups of $n + 1$ siblings to a vertex S_n .




first idea: look at small examples and find a bijection


S_n

\leftrightarrow



T_n





\leftrightarrow




\leftrightarrow




\leftrightarrow



\leftrightarrow



first idea: look at small examples and find a bijection



second idea: follow Read, Gessel & Li and count

combinatorial species (Joyal 1981)

a combinatorial species is ...

Definition. ... a gadget to enumerate symmetry classes of labelled objects. Given a set of *labels*, a species 'produces' the set of *(labelled) structures*.

- ▶ *Par*, set partitions

$$Par[a, 1, \heartsuit] = \{a1\heartsuit, a1|\heartsuit, a\heartsuit|1, 1\heartsuit|a, a|1|\heartsuit\}$$

a combinatorial species is ...

Definition. ... a gadget to enumerate symmetry classes of labelled objects. Given a set of *labels*, a species 'produces' the set of (*labelled*) *structures*.

- ▶ *Par*, set partitions

$$Par[a, 1, \heartsuit] = \{a1\heartsuit, a1|\heartsuit, a\heartsuit|1, 1\heartsuit|a, a|1|\heartsuit\}$$

- ▶ \mathcal{G} , (vertex labelled simple) graphs

a combinatorial species is ...

Definition. ... a gadget to enumerate symmetry classes of labelled objects. Given a set of *labels*, a species 'produces' the set of (*labelled*) *structures*.

- ▶ *Par*, set partitions

$$Par[a, 1, \heartsuit] = \{a1\heartsuit, a1|\heartsuit, a\heartsuit|1, 1\heartsuit|a, a|1|\heartsuit\}$$

- ▶ \mathcal{G} , (vertex labelled simple) graphs
- ▶ E , sets, and E_n , sets of size n

$$E[a, 1, \heartsuit] = \{\{a, 1, \heartsuit\}\} \quad E_2[a, 1, \heartsuit] = \{\}$$

a combinatorial species is ...

Definition. ... a functor F from the category of finite sets with bijections to the category of finite sets with maps:

- ▶ for a finite set of *labels* U ,
 $F[U]$ is the finite set of *F -structures on U* ,

a combinatorial species is ...

Definition. ... a functor F from the category of finite sets with bijections to the category of finite sets with maps:

- ▶ for a finite set of *labels* U ,
 $F[U]$ is the finite set of *F -structures on U* ,
- ▶ for a bijection $\sigma : U \rightarrow V$,
 $F[\sigma] : F[U] \rightarrow F[V]$ is the *relabelling* of the structures,

such that

$$F[id_U] = id_{F[U]}$$

and, for any two bijections $\sigma : U \rightarrow V$ and $\tau : V \rightarrow W$,

$$F[\tau \circ \sigma] = F[\tau] \circ F[\sigma].$$

a combinatorial species is ...

Definition. ... a functor F from the category of finite sets with bijections to the category of finite sets with maps:

- ▶ for a finite set of *labels* U ,
 $F[U]$ is the finite set of *F -structures on U* ,
- ▶ for a bijection $\sigma : U \rightarrow V$,
 $F[\sigma] : F[U] \rightarrow F[V]$ is the *relabelling* of the structures,

such that

$$F[id_U] = id_{F[U]}$$

and, for any two bijections $\sigma : U \rightarrow V$ and $\tau : V \rightarrow W$,

$$F[\tau \circ \sigma] = F[\tau] \circ F[\sigma].$$

usually, the definition of relabelling is clear from context!

a combinatorial species is ...

Definition. ... a functor F from the category of finite sets with bijections to the category of finite sets with maps:

- ▶ for a finite set of *labels* U ,
 $F[U]$ is the finite set of *F -structures on U* ,
- ▶ for a bijection $\sigma : U \rightarrow V$,
 $F[\sigma] : F[U] \rightarrow F[V]$ is the *relabelling* of the structures,

such that

$$F[id_U] = id_{F[U]}$$

and, for any two bijections $\sigma : U \rightarrow V$ and $\tau : V \rightarrow W$,

$$F[\tau \circ \sigma] = F[\tau] \circ F[\sigma].$$

$$Par[a, 1, \heartsuit] = \{a1\heartsuit, a1|\heartsuit, a\heartsuit|1, 1\heartsuit|a, a|1|\heartsuit\}$$

$$Par\left[\begin{pmatrix} a & 1 & \heartsuit \\ x & y & z \end{pmatrix}\right](a1|\heartsuit) = xy|z$$

isomorphism of structures

Definition. Structures $s \in F[U]$ and $t \in F[V]$ are *isomorphic* if t is obtained from s by relabelling:

$$t = F[\sigma](s) \quad \text{for a bijection } \sigma : U \rightarrow V$$

$(a1|\heartsuit) \in \text{Par}[a, 1, \heartsuit]$ and $(xy|z) \in \text{Par}[x, y, z]$ are isomorphic, because

$$\text{Par}\left[\begin{pmatrix} a & 1 & \heartsuit \\ x & y & z \end{pmatrix}\right](a1|\heartsuit) = xy|z.$$

computing with species

Definition.

$$(F + G)[U] = F[U] + G[U]$$

sum

computing with species


Definition.

$$(F + G)[U] = F[U] + G[U]$$

sum

$$(F \cdot G)[U] = \sum_{V+W=U} F[V] \times G[W]$$


product



computing with species

Definition.

$$(F + G)[U] = F[U] + G[U] \quad \text{sum}$$
$$(F \cdot G)[U] = \sum_{V+W=U} F[V] \times G[W] \quad \text{product}$$
$$(F \circ G)[U] = \sum_{\pi \in \text{Par}[U]} F[\pi] \times \prod_{B \in \pi} G[B] \quad \text{composition}$$



computing with species

Definition.

$$\begin{aligned}(F + G)[U] &= F[U] + G[U] && \text{sum} \\(F \cdot G)[U] &= \sum_{V+W=U} F[V] \times G[W] && \text{product} \\(F \circ G)[U] &= \sum_{\pi \in \text{Par}[U]} F[\pi] \times \prod_{B \in \pi} G[B] && \text{composition}\end{aligned}$$

Definition.

$$\begin{aligned}F(x) &= \sum_{n \geq 0} |F[n]| \frac{x^n}{n!} \\ \tilde{F}(x) &= \sum_{n \geq 0} \# \text{isomorphism types of } F[n] x^n\end{aligned}$$

computing with species

Definition.

$$\begin{aligned}(F + G)[U] &= F[U] + G[U] && \text{sum} \\(F \cdot G)[U] &= \sum_{V+W=U} F[V] \times G[W] && \text{product} \\(F \circ G)[U] &= \sum_{\pi \in \text{Par}[U]} F[\pi] \times \prod_{B \in \pi} G[B] && \text{composition}\end{aligned}$$

Definition.

$$\begin{aligned}F(x) &= \sum_{n \geq 0} |F[n]| \frac{x^n}{n!} \\ \widetilde{F}(x) &= \sum_{n \geq 0} \# \text{isomorphism types of } F[n] x^n\end{aligned}$$

Proposition.

$$\begin{aligned}(F + G)(x) &= F(x) + G(x), & (F \cdot G)(x) &= F(x)G(x), & (F \circ G)(x) &= F(G(x)) \\ \widetilde{(F + G)}(x) &= \widetilde{F}(x) + \widetilde{G}(x), & \widetilde{(F \cdot G)}(x) &= \widetilde{F}(x)\widetilde{G}(x), & \widetilde{(F \circ G)}(x) &= \widetilde{F}(\widetilde{G}(x))\end{aligned}$$

computing with species

Definition. $Z_F(x_1, x_2, \dots) = \sum_{n \geq 0} \frac{1}{n!} \sum_{\sigma \in \mathfrak{S}_n} \text{fix } F[\sigma] x_1^{\sigma_1} x_2^{\sigma_2} \dots,$

where

$\text{fix } F[\sigma] = \# \text{fixed points of } F[\sigma],$

$\sigma_k = \#k\text{-cycles of } \sigma,$

is the *cycle index series* or *Frobenius character* of F .

Example. $E(x) = \exp(x), \quad \tilde{E}(x) = \frac{1}{1-x}, \quad Z_E = \exp \sum_{n > 0} \frac{x_n}{n}$

computing with species

Definition. $Z_F(x_1, x_2, \dots) = \sum_{n \geq 0} \frac{1}{n!} \sum_{\sigma \in \mathfrak{S}_n} \text{fix } F[\sigma] x_1^{\sigma_1} x_2^{\sigma_2} \dots,$

where

$\text{fix } F[\sigma] = \# \text{fixed points of } F[\sigma],$

$\sigma_k = \#k\text{-cycles of } \sigma,$

is the *cycle index series* or *Frobenius character* of F .

Example. $E(x) = \exp(x), \quad \tilde{E}(x) = \frac{1}{1-x}, \quad Z_E = \exp \sum_{n > 0} \frac{x_n}{n}$

Fact. $F(x) = Z_F(x, 0, \dots), \quad \tilde{F}(x) = Z_F(x, x^2, \dots)$

Theorem. $\widetilde{F \circ G}(x) = Z_F(\tilde{G}(x), \tilde{G}(x^2), \dots)$

molecular decompositions and virtual species

Definition. F is *molecular* if it is not a non-trivial sum.

Fact. F is molecular if any two F structures are isomorphic.

$F \cdot G$ is molecular if F and G are molecular.

The molecular decomposition is unique.

$$\begin{aligned} \mathcal{G} = & 1 + X + 2E_2 + (2E_3 + 2XE_2) \\ & + (2E_4 + 2E_2^2 + 2XE_3 + 2P_4 + 2X^2E_2 + E_2(X^2)) + \dots \end{aligned}$$

molecular decompositions and virtual species

Definition. F is *molecular* if it is not a non-trivial sum.

Fact. F is molecular if any two F structures are isomorphic.

$F \cdot G$ is molecular if F and G are molecular.

The molecular decomposition is unique.

$$\begin{aligned} \mathcal{G} = & 1 + X + 2E_2 + (2E_3 + 2XE_2) \\ & + (2E_4 + 2E_2^2 + 2XE_3 + 2P_4 + 2X^2E_2 + E_2(X^2)) + \dots \end{aligned}$$

Definition. A *virtual species* is a formal sum of molecular species.

Fact. F has a compositional inverse if $F_0 = 0$ and $F_1 \neq 0$.

$$E_{\geq 1}^{(-1)} = X + (-E_2) + (-E_3 + XE_2) + (-E_4 + P_4 + XE_3 - X^2E_2) + \dots$$

Read's idea

- ▶ \mathcal{G} ... graphs
- ▶ \mathcal{M} ... co-mating graphs

Theorem (Read 1989). $\mathcal{M} \circ E_{\geq 1} = \mathcal{G}$.

Read's idea

- ▶ \mathcal{G} ... graphs
- ▶ \mathcal{M} ... co-mating graphs

Theorem (Read 1989). $\mathcal{M} \circ E_{\geq 1} = \mathcal{G}$.

Proof. $(F \circ G)[U] = \sum_{\pi \in \text{Par}[U]} F[\pi] \times \prod_{B \in \pi} G[B]$.

$$\underbrace{\begin{array}{c} \textcircled{1} \text{---} \textcircled{2} \\ \text{---} \textcircled{3} \\ \text{---} \textcircled{4} \end{array}}_{\in \mathcal{G}[1,2,3,4]} = \left(\underbrace{\{\{1\}, \{2\}, \{3,4\}\}}_{\in \text{Par}[1,2,3,4]}, \underbrace{\textcircled{1} \text{---} \textcircled{2} \text{---} \textcircled{34}}_{\in \mathcal{M}[\{1\}, \{2\}, \{3,4\}]}, (\{1\}, \{2\}, \{3,4\}) \right)$$



Read's idea

- ▶ \mathcal{G} ... graphs
- ▶ \mathcal{M} ... co-mating graphs

Theorem (Read 1989). $\mathcal{M} \circ E_{\geq 1} = \mathcal{G}$.

Proof. $(F \circ G)[U] = \sum_{\pi \in \text{Par}[U]} F[\pi] \times \prod_{B \in \pi} G[B]$.

$$\underbrace{\begin{array}{c} \textcircled{1} \text{---} \textcircled{2} \\ \text{---} \textcircled{4} \\ \text{---} \textcircled{3} \end{array}}_{\in \mathcal{G}[1,2,3,4]} = \left(\underbrace{\{\{1\}, \{2\}, \{3,4\}\}}_{\in \text{Par}[1,2,3,4]}, \underbrace{\textcircled{1} \text{---} \textcircled{2} \text{---} \textcircled{34}}_{\in \mathcal{M}[\{1\}, \{2\}, \{3,4\}]}, (\{1\}, \{2\}, \{3,4\}) \right)$$

□

Corollary (Read 1989). $\mathcal{M} = \mathcal{G} \circ E_{\geq 1}^{(-1)}$.

Gessel & Li's idea: a tuft decomposition

- ▶ $X + \mathcal{T} := X \cdot E(Y)$... a singleton vertex or a tuft
- ▶ $\mathcal{G}_{\neq 2}(X, Y)$... graphs with leaves Y


Theorem (Gessel & Li 2011). $\mathcal{G}_{\neq 2}(X, X + Y) = \mathcal{G}(X + \mathcal{T}) - XY - E_2.$

Gessel & Li's idea: a tuft decomposition

- ▶ $X + \mathcal{T} := X \cdot E(Y)$... a singleton vertex or a tuft
- ▶ $\mathcal{G}_{\neq 2}(X, Y)$... graphs with leaves Y

Theorem (Gessel & Li 2011). $\mathcal{G}_{\neq 2}(X, X + Y) = \mathcal{G}(X + \mathcal{T}) - XY - E_2$.

Proof.



new idea: a patch decomposition

- ▶ $\mathcal{P}(X, Y, Z) := (1 + X + E_{\geq 2}(Z)) \cdot (E \circ \mathcal{T}) - 1$... a *patch*
- ▶ $\mathcal{G}_{\neq 2}(X, Y, Z)$... graphs with leaves Y , siblings Z

Theorem (Fürnsinn & Gangl & R. 2024).

$$\mathcal{G}_{\neq 2}(X, X + Y, Z) = \mathcal{M} \circ \mathcal{P} - XY - E_2(Z).$$


new idea: a patch decomposition

- ▶ $\mathcal{P}(X, Y, Z) := (1 + X + E_{\geq 2}(Z)) \cdot (E \circ \mathcal{T}) - 1 \dots$ a *patch*
- ▶ $\mathcal{G}_{\neq 2}(X, Y, Z) \dots$ graphs with leaves Y , siblings Z

Theorem (Fürsinn & Gangl & R. 2024).

$$\mathcal{G}_{\neq 2}(X, X + Y, Z) = \mathcal{M} \circ \mathcal{P} - XY - E_2(Z).$$

Proof.




new idea: a patch decomposition

- ▶ $\mathcal{P}^s(X, Y, Z) := (1 + X + E_{2 \leq \bullet < 2+s}(Z)) \cdot (E \circ T) - 1 \dots$ a *patch*
- ▶ $\mathcal{G}_{\neq 2}^s(X, Y, Z) \dots$ graphs with leaves Y , siblings Z ,
sibling number $\leq s$

Theorem (Fürnsinn & Gangl & R. 2024).

$$\mathcal{G}_{\neq 2}^s(X, X + Y, Z) = \mathcal{M} \circ \mathcal{P}^s - XY - [s > 0]E_2(Z).$$

Proof.



□

harvest

Lemma. Let

$$\mathcal{V}^{s,t} := \left(1 + X + E_{2 \leq \bullet < 2+s}(Z)\right) \cdot E\left(X \cdot E_{<1+t}(Y)E(-X) - 1\right) - 1.$$

Then $\mathcal{G}_{\neq 2}^{s,t}(X, Y, Z) = \mathcal{M} \circ \mathcal{V}^{s,t} + X^2 - [t > 0]XY - [s > 0]E_2(Z)$.

Proof.

$Y := Y - X$ in the patch decomposition, and similar tricks. □

harvest

Lemma. Let

$$\mathcal{V}^{s,t} := (1 + X + E_{2 \leq \bullet < 2+s}(Z)) \cdot E(X \cdot E_{< 1+t}(Y)E(-X) - 1) - 1.$$

Then $\mathcal{G}_{\neq 2}^{s,t}(X, Y, Z) = \mathcal{M} \circ \mathcal{V}^{s,t} + X^2 - [t > 0]XY - [s > 0]E_2(Z)$.

Proof.

$Y := Y - X$ in the patch decomposition, and similar tricks. □

Lemma. $\widetilde{\mathcal{V}}^{s,t}(x) := \widetilde{\mathcal{V}}^{s,t}(x, x, x) = \frac{(1-x^{s+2})(1-x^{t+2})}{1-x} - 1$.

Corollary. $\widetilde{\mathcal{G}}_{\neq 2}^{s,t}(x, x, x)$ is symmetric in s and t .

Proof.

$$\widetilde{\mathcal{M}} \circ \widetilde{\mathcal{V}}^{s,t} = Z_{\mathcal{M}}(\widetilde{\mathcal{V}}^{s,t}(x), \widetilde{\mathcal{V}}^{s,t}(x^2), \dots).$$

□

new idea: a refined patch decomposition

Definition. The *reduction* of a graph other than K_2 is obtained by repeatedly removing leaves and contracting each group of siblings to a single vertex.

- ▶ $\mathcal{G}_{\neq 2}^{R,s}(X, Y, Z)$... graphs with leaves Y , siblings Z , sibling number $\leq s$, that reduce to R
- ▶ \mathcal{M}^R ... co-mating graphs that reduce to R

new idea: a refined patch decomposition

Definition. The *reduction* of a graph other than K_2 is obtained by repeatedly removing leaves and contracting each group of siblings to a single vertex.

- ▶ $\mathcal{G}_{\neq 2}^{R,s}(X, Y, Z)$... graphs with leaves Y , siblings Z , sibling number $\leq s$, that reduce to R
- ▶ \mathcal{M}^R ... co-mating graphs that reduce to R

Theorem (Fürnsinn & Gangl & R. 2024).

$$\mathcal{G}_{\neq 2}^{R,s}(X, X + Y, Z) = \begin{cases} \mathcal{M}^\bullet \circ \mathcal{P}^s - XY - [s > 0]E_2(Z) & R = \bullet \\ \mathcal{M}^R \circ \mathcal{P}^s & \text{otherwise.} \end{cases}$$

Theorem (Fürnsinn & Gangl & R. 2024). On the set of graphs that reduce to R ,

$$(s, t) \sim (t, s)$$

questions

- ▶ find a bijection!

questions

- ▶ find a bijection!
- ▶ find any bijection!

questions

- ▶ find a bijection!
- ▶ find any bijection!
- ▶ do tropical species exist?