

# Stochastic diffusion model on the unit circle

---


Zoé Varin

February 5th, 2025


Joint work with Jean-François Marckert

## Definition of the model

State space  $\mathcal{C} = \mathbb{R}/\mathbb{Z}$ .  $m_0, \dots, m_n$  with  $\sum m_i < 1$ .



## Definition of the model




State space  $\mathcal{C} = \mathbb{R}/\mathbb{Z}$ .  $m_0, \dots, m_n$  with  $\sum m_i < 1$ .

At every step  $k$ :

- arrival of a drop of mass  $m_k$  at  $u_k \sim \mathcal{U}(\mathcal{C})$

## Definition of the model




State space  $\mathcal{C} = \mathbb{R}/\mathbb{Z}$ .  $m_0, \dots, m_n$  with  $\sum m_i < 1$ .

At every step  $k$ :

- arrival of a drop of mass  $m_k$  at  $u_k \sim \mathcal{U}(\mathcal{C})$
- continuous spreading of the drop (so that the new covered area has size  $m_k$ )

## Definition of the model




State space  $\mathcal{C} = \mathbb{R}/\mathbb{Z}$ .  $m_0, \dots, m_n$  with  $\sum m_i < 1$ .

At every step  $k$ :

- arrival of a drop of mass  $m_k$  at  $u_k \sim \mathcal{U}(\mathcal{C})$
- continuous spreading of the drop (so that the new covered area has size  $m_k$ )

## Definition of the model




State space  $\mathcal{C} = \mathbb{R}/\mathbb{Z}$ .  $m_0, \dots, m_n$  with  $\sum m_i < 1$ .

At every step  $k$ :

- arrival of a drop of mass  $m_k$  at  $u_k \sim \mathcal{U}(\mathcal{C})$
- continuous spreading of the drop (so that the new covered area has size  $m_k$ )

## Definition of the model




State space  $\mathcal{C} = \mathbb{R}/\mathbb{Z}$ .  $m_0, \dots, m_n$  with  $\sum m_i < 1$ .

At every step  $k$ :

- arrival of a drop of mass  $m_k$  at  $u_k \sim \mathcal{U}(\mathcal{C})$
- continuous spreading of the drop (so that the new covered area has size  $m_k$ )

## Definition of the model




State space  $\mathcal{C} = \mathbb{R}/\mathbb{Z}$ .  $m_0, \dots, m_n$  with  $\sum m_i < 1$ .

At every step  $k$ :

- arrival of a drop of mass  $m_k$  at  $u_k \sim \mathcal{U}(\mathcal{C})$
- continuous spreading of the drop (so that the new covered area has size  $m_k$ )



## Definition of the model




State space  $\mathcal{C} = \mathbb{R}/\mathbb{Z}$ .  $m_0, \dots, m_n$  with  $\sum m_i < 1$ .

At every step  $k$ :

- arrival of a drop of mass  $m_k$  at  $u_k \sim \mathcal{U}(\mathcal{C})$
- continuous spreading of the drop (so that the new covered area has size  $m_k$ )

## Definition of the model




State space  $\mathcal{C} = \mathbb{R}/\mathbb{Z}$ .  $m_0, \dots, m_n$  with  $\sum m_i < 1$ .

At every step  $k$ :

- arrival of a drop of mass  $m_k$  at  $u_k \sim \mathcal{U}(\mathcal{C})$
- continuous spreading of the drop (so that the new covered area has size  $m_k$ )

## Definition of the model




State space  $\mathcal{C} = \mathbb{R}/\mathbb{Z}$ .  $m_0, \dots, m_n$  with  $\sum m_i < 1$ .

At every step  $k$ :

- arrival of a drop of mass  $m_k$  at  $u_k \sim \mathcal{U}(\mathcal{C})$
- continuous spreading of the drop (so that the new covered area has size  $m_k$ )

## Definition of the model




State space  $\mathcal{C} = \mathbb{R}/\mathbb{Z}$ .  $m_0, \dots, m_n$  with  $\sum m_i < 1$ .

At every step  $k$ :

- arrival of a drop of mass  $m_k$  at  $u_k \sim \mathcal{U}(\mathcal{C})$
- continuous spreading of the drop (so that the new covered area has size  $m_k$ )

# Definition of the model



State space  $\mathcal{C} = \mathbb{R}/\mathbb{Z}$ .  $m_0, \dots, m_n$  with  $\sum m_i < 1$ .


At every step  $k$ :

- arrival of a drop of mass  $m_k$  at  $u_k \sim \mathcal{U}(\mathcal{C})$
- continuous spreading of the drop (so that the new covered area has size  $m_k$ )

Configuration at time  $k$  (i.e. after drop  $k - 1$  has been spread):

- **occupied space**  $O^{(k)}$  of size  $\text{Leb}(O^{(k)}) = \sum_{i=0}^{k-1} m_i$
- **free space**  $F^{(k)} = \mathcal{C} \setminus O^{(k)}$

## Definition of the model



State space  $\mathcal{C} = \mathbb{R}/\mathbb{Z}$ .  $m_0, \dots, m_n$  with  $\sum m_i < 1$ .

At every step  $k$ :


- arrival of a drop of mass  $m_k$  at  $u_k \sim \mathcal{U}(\mathcal{C})$
- continuous spreading of the drop (so that the new covered area has size  $m_k$ )

Configuration at time  $k$  (i.e. after drop  $k - 1$  has been spread):

- **occupied space**  $O^{(k)}$  of size  $\text{Leb}(O^{(k)}) = \sum_{i=0}^{k-1} m_i$
- **free space**  $F^{(k)} = \mathcal{C} \setminus O^{(k)}$
- more precisely,  $N^{(k)}$  **blocks** of each type

$\Rightarrow (O_i^{(k)}, F_i^{(k)})_{1 \leq i \leq N^{(k)}}$  ordered around the circle with  $0 \in O_1^{(k)} \cup F_1^{(k)}$

## Definition of the model



State space  $\mathcal{C} = \mathbb{R}/\mathbb{Z}$ .  $m_0, \dots, m_n$  with  $\sum m_i < 1$ .

At every step  $k$ :

- arrival of a drop of mass  $m_k$  at  $u_k \sim \mathcal{U}(\mathcal{C})$
- continuous spreading of the drop (so that the new covered area has size  $m_k$ )

Configuration at time  $k$  (i.e. after drop  $k - 1$  has been spread):

- **occupied space**  $O^{(k)}$  of size  $\text{Leb}(O^{(k)}) = \sum_{i=0}^{k-1} m_i$
- **free space**  $F^{(k)} = \mathcal{C} \setminus O^{(k)}$
- more precisely,  $N^{(k)}$  **blocks** of each type

$\Rightarrow (O_i^{(k)}, F_i^{(k)})_{1 \leq i \leq N^{(k)}}$  ordered around the circle with  $0 \in O_1^{(k)} \cup F_1^{(k)}$

# Examples of spreading policies

- Right diffusion at constant speed:  $\overrightarrow{O^{(k)}}, \overrightarrow{F^{(k)}}$




(studied by Bertoin, Miermont [BM06])



# Examples of spreading policies

- **Right diffusion at constant speed:**  $\overrightarrow{O^{(k)}}, \overrightarrow{F^{(k)}}$



(studied by Bertoin, Miermont [BM06])

- Diffusion to the closest side (with or without constant reevaluation)
- Range of a brownian path
- short-sighted jam spreader
- for any ball, pick at random some spreading policy
- ...

# What is a valid spreading policy ?

As a continuous process: local and continuous diffusion



# What is a valid spreading policy ?

As a continuous process: local and continuous diffusion



Hypotheses:

- $dl$  and  $dr$  only depend on what is inside the current component of  $u_k$  (one of the  $O_i^{(k+\varepsilon)}$ )
- invariance by translation of the process

## A universality result

We consider  $\frac{\sigma \cdot |F^{(k)}|}{R} = \left( \frac{|F_{\sigma_i}^{(k)}|}{R} \right)_{1 \leq i \leq N^{(k)}}$ , for  $\sigma \sim \mathcal{U}(\mathfrak{S}_{N^{(k)}})$ . Let  $R = 1 - \sum_{i=0}^{k-1} m_i$ .

### Theorem


*Independently of the diffusion policy,*

- **Number of blocks:**  $N^{(k)} \stackrel{(d)}{=} 1 + \text{Binomial}(k-1, R)$
- **Lengths of the free blocks:**  $\frac{\sigma \cdot |F^{(k)}|}{R} \sim \text{Dirichlet}(N^{(k)}; 1, \dots, 1)$
- **Lengths of the occupied blocks:** a formula for  $\mathbb{P}(|O^{(k)}| = (M_0, \dots, M_{b-1}))$

# A universality result

We consider  $\frac{\sigma \cdot |F^{(k)}|}{R} = \left( \frac{|F_{\sigma_i}^{(k)}|}{R} \right)_{1 \leq i \leq N^{(k)}}$ , for  $\sigma \sim \mathcal{U}(\mathcal{S})$


$$u_1, u_2, u_3 \sim \mathcal{U}([0, 1])$$



## Theorem

*Independently of the diffusion policy,*

- **Number of blocks:**  $N^{(k)} \stackrel{(d)}{=} 1 + \text{Binomial}(k, \frac{1}{2})$
- **Lengths of the free blocks:**  $\frac{\sigma \cdot |F^{(k)}|}{R} \sim \text{Dirichlet}(N^{(k)}, \mathbf{1}, \dots, \mathbf{1})$
- **Lengths of the occupied blocks:** a formula for  $\mathbb{P}(|O^{(k)}| = (M_0, \dots, M_{b-1}))$



## A universality result

We consider  $\frac{\sigma \cdot |F^{(k)}|}{R} = \left( \frac{|F_{\sigma_i}^{(k)}|}{R} \right)_{1 \leq i \leq N^{(k)}}$ , for  $\sigma \sim \mathcal{U}(\mathfrak{S}_{N^{(k)}})$ . Let  $R = 1 - \sum_{i=0}^{k-1} m_i$ .

### Theorem


*Independently of the diffusion policy,*

- **Number of blocks:**  $N^{(k)} \stackrel{(d)}{=} 1 + \text{Binomial}(k-1, R)$
- **Lengths of the free blocks:**  $\frac{\sigma \cdot |F^{(k)}|}{R} \sim \text{Dirichlet}(N^{(k)}; 1, \dots, 1)$
- **Lengths of the occupied blocks:** a formula for  $\mathbb{P}(|O^{(k)}| = (M_0, \dots, M_{b-1}))$


# Some background on the continuous and discrete parking models

## Discrete parking

- introduced by Konheim, Weiss [KW66], studied by Knuth [Knu73]




- asymptotic behavior studied by Chassaing, Louchard [CL02]




# Some background on the continuous and discrete parking models

## Discrete parking

- introduced by Konheim, Weiss [KW66], studied by Knuth [Knu73]




- asymptotic behavior studied by Chassaing, Louchard [CL02]



## Additive coalescent

studied by Aldous, Pitman [AP98], Chassaing, Louchard,...







# Some background on the continuous and discrete parking models

## Discrete parking

- introduced by Konheim, Weiss [KW66], studied by Knuth [Knu73]



- asymptotic behavior studied by Chassaing, Louchard [CL02]




## Generalized parking

- Parking on  $\mathbb{Z}$  (Przykucki, Roberts, Scott [PRS23])
- Parking on (random) trees (Contat et. al.)
- Bilateral parking procedures (Nadeau), Golf model [Var24]

## Additive coalescent


studied by Aldous, Pitman [AP98], Chassaing, Louchard,...




# Some background on the continuous and discrete parking models

## Discrete parking

- introduced by Konheim, Weiss [KW66], studied by Knuth [Knu73]



- asymptotic behavior studied by Chassaing, Louchard [CL02]




**Continuous version of the classical parking**  
Caravans (Bertoin, Miermont [BM06])

## Generalized parking

- Parking on  $\mathbb{Z}$  (Przykucki, Roberts, Scott [PRS23])
- Parking on (random) trees (Contat et. al.)
- Bilateral parking procedures (Nadeau), Golf model [Var24]

## Additive coalescent


studied by Aldous, Pitman [AP98], Chassaing, Louchard,...




# Some background on the continuous and discrete parking models

## Discrete parking

- introduced by Konheim, Weiss [KW66], studied by Knuth [Knu73]



- asymptotic behavior studied by Chassaing, Louchard [CL02]




## Generalized parking

- Parking on  $\mathbb{Z}$  (Przykucki, Roberts, Scott [PRS23])
- Parking on (random) trees (Contat et. al.)
- Bilateral parking procedures (Nadeau), Golf model [Var24]

## Additive coalescent

studied by Aldous, Pitman [AP98], Chassaing, Louchard,...



**Continuous version of the classical parking**  
Caravans (Bertoin, Miermont [BM06])



## A universality result

We consider  $\frac{\sigma \cdot |F^{(k)}|}{R} = \left( \frac{|F_{\sigma_i}^{(k)}|}{R} \right)_{1 \leq i \leq N^{(k)}}$ , for  $\sigma \sim \mathcal{U}(\mathfrak{S}_{N^{(k)}})$ . Let  $R = 1 - \sum_{i=0}^{k-1} m_i$ .

### Theorem

*Independently of the diffusion policy,*

- **Number of blocks:**  $N^{(k)} \stackrel{(d)}{=} 1 + \text{Binomial}(k-1, R)$
- **Lengths of the free blocks:**  $\frac{\sigma \cdot |F^{(k)}|}{R} \sim \text{Dirichlet}(N^{(k)}; 1, \dots, 1)$
- **Lengths of the occupied blocks:** a formula for  $\mathbb{P}(|O^{(k)}| = (M_0, \dots, M_{b-1}))$

# Main principle

Consider 4 uniform points on  $[0, 1]$ .

Conditional on :



Then:





4 uniform points in 



3 uniform points in 


# Intuition on the proof

Peaks representation:




# Intuition on the proof

Peaks representation:



# Intuition on the proof


Peaks representation:





# Intuition on the proof

## Peaks representation:




allow to see that, at every step:

- the positions of the peaks are uniform on the smaller cycle  $\mathcal{C}_R$  of size  $R = 1 - \sum m_i$
- the distributions of the peaks **number**, **heights** and **positions** do not depend on the diffusion policy

# Intuition on the proof

## Peaks representation:




allow to see that, at every step:

- the positions of the peaks are uniform on the smaller cycle  $\mathcal{C}_R$  of size  $R = 1 - \sum m_i$
- the distributions of the peaks **number**, **heights** and **positions** do not depend on the diffusion policy

# Intuition on the proof

## Peaks representation:



allow to see that, at every step:

- the positions of the peaks are uniform on the smaller cycle  $\mathcal{C}_R$  of size  $R = 1 - \sum m_i$
- the distributions of the peaks **number**, **heights** and **positions** do not depend on the diffusion policy

# Distribution of the lengths of the free blocks

Reminder:  $|F| = (|F_i|)_{1 \leq i \leq N^{(k)}}$  and  $\frac{\sigma \cdot |F|}{R} = \left( \frac{|F_{\sigma_i}|}{R} \right)_{1 \leq i \leq N^{(k)}}$ ,  
for any  $\sigma \in \mathfrak{S}_{N^{(k)}}$ .



## Theorem (Distribution of $|F|$ conditional on $N^{(k)}$ )

Recall that  $R = 1 - \sum m_i$ .

- **unbiased version:** if  $\sigma \sim \mathcal{U}(\mathfrak{S}_{N^{(k)}})$ ,  $\frac{\sigma \cdot |F|}{R} \sim \text{Dirichlet}(N^{(k)}; 1, \dots, 1)$

# Distribution of the lengths of the free blocks

Reminder:  $|F| = (|F_i|)_{1 \leq i \leq N^{(k)}}$  and  $\frac{\sigma \cdot |F|}{R} = \left( \frac{|F_{\sigma_i}|}{R} \right)_{1 \leq i \leq N^{(k)}}$ ,  
for any  $\sigma \in \mathfrak{S}_{N^{(k)}}$ .



## Theorem (Distribution of $|F|$ conditional on $N^{(k)}$ )


Recall that  $R = 1 - \sum m_i$ .

- **unbiased version:** if  $\sigma \sim \mathcal{U}(\mathfrak{S}_{N^{(k)}})$ ,  $\frac{\sigma \cdot |F|}{R} \sim \text{Dirichlet}(N^{(k)}; 1, \dots, 1)$
- **biased version:**

$$\frac{(|F_1|, \dots, |F_{N^{(k)}}|)}{R} \sim \begin{cases} \text{Dirichlet}(N^{(k)}; 1, \dots, 1) & \text{with probability } 1 - R \\ \text{Dirichlet}(N^{(k)}; 2, 1, \dots, 1) & \text{with probability } R \end{cases}$$

# Intuition on the proof (bis)

## Peaks representation:




allow to see that, at every step:

- the positions of the peaks are uniform on the smaller cycle  $\mathcal{C}_R$  of size  $R = 1 - \sum m_i$
- the distributions of the peaks **number**, **heights** and **positions** do not depend on the diffusion policy

# Intuition on the proof (bis)

## Peaks representation:



allow to see that, at every step:

- the positions of the peaks are uniform on the smaller cycle  $\mathcal{C}_R$  of size  $R = 1 - \sum m_i$
- the distributions of the peaks **number**, **heights** and **positions** do not depend on the diffusion policy
- even more surprisingly, for the peaks **number** and **positions**: do not depend on which peak is extended by the diffusion

# Distribution of the number of blocks $N^{(k)}$

## Theorem

$$\mathcal{L}\left(|F^{(k)}| \mid (m_0, \dots, m_{k-1})\right) = \mathcal{L}\left(\overrightarrow{|F^{(k)}|} \mid (m_0, \dots, m_{k-1})\right)$$



## Distribution of the number of blocks $N^{(k)}$

### Theorem

$$\mathcal{L}\left(|F^{(k)}| \mid (m_0, \dots, m_{k-1})\right) = \mathcal{L}\left(|\vec{F}^{(k)}| \mid (\sum m_i, 0, \dots, 0)\right)$$

$$\mathcal{L}\left(|F^{(k)}| \mid (m_0, \dots, m_{k-1})\right) = \mathcal{L}\left(|\vec{F}^{(k)}| \mid (m_0, \dots, m_{k-1})\right)$$

# Distribution of the number of blocks $N^{(k)}$

## Theorem

$$\mathcal{L}\left(|F^{(k)}| \mid (m_0, \dots, m_{k-1})\right) = \mathcal{L}\left(|\overrightarrow{F^{(k)}}| \mid (\sum m_i, 0, \dots, 0)\right)$$

## Theorem (Distribution of $N^{(k)}$ )

Let  $B(k-1, R) \sim \text{Binomial}(k-1, R)$ , then

$$N^{(k)} \stackrel{(d)}{=} 1 + B(k-1, R)$$




$$N^{(k)} = 1 + \sum_{j=1}^{k-1} \mathbb{1}_{u_k \notin \text{blue}}$$

# Distribution of the occupied blocks

## One block case:

$$\mathbb{P}\left(N^{(k)} = 1\right) = \left(\sum m_i\right)^{k-1} =: Q\left(\sum m_i, k\right)$$

and, conditional on  $N^{(k)} = 1$ ,  $O^{(k)}$  is reduced to an interval  $[A, A + \sum m_i]$  with  $A$  uniform on  $\mathcal{C}$




# Distribution of the occupied blocks

## One block case:

$$\mathbb{P}\left(N^{(k)} = 1\right) = \left(\sum m_i\right)^{k-1} =: Q\left(\sum m_i, k\right)$$

and, conditional on  $N^{(k)} = 1$ ,  $O^{(k)}$  is reduced to an interval  $[A, A + \sum m_i]$  with  $A$  uniform on  $\mathcal{C}$



## General case:

### Theorem

$$\mathbb{P}\left(|O^{(k)}| = (M_0, \dots, M_{b-1})\right) = T(M_0, \dots, M_{b-1}) \sum_{P \in \mathcal{P}(k, b)} \left[ \prod_{\ell=0}^{b-1} Q(M_\ell, |P_\ell|) \mathbf{1}_{\sum_{i \in P_\ell} m_i = M_\ell} \right]$$

where

-  $\mathcal{P}(k, b)$  is the set of partitions  $P = (P_0, \dots, P_{b-1})$  of  $\{1, \dots, k-1\}$  into  $b$  non empty parts,

-  $T(M_0, \dots, M_{b-1}) = M_0 \frac{(1 - \sum M_\ell)^{b-1}}{(b-1)!} + \frac{(1 - \sum M_\ell)^b}{b!}$ .

# Summary of universality results

## Theorem

For any continuous model with valid spreading policy, the following distributions are **explicit** and **independent of the spreading policy**:

- With  $k$  fixed:

- $\mathcal{L}(O^{(k)}, F^{(k)})$

- $\mathcal{L}\left(|F^{(k)}| \mid (m_0, \dots, m_{k-1})\right) = \mathcal{L}\left(|\vec{F}^{(k)}| \mid (\sum m_i, 0, \dots, 0)\right)$

## Theorem

For any continuous model with valid spreading policy, the following distributions are **explicit** and **independent of the spreading policy**:

- With  $k$  fixed:

- $\mathcal{L}(O^{(k)}, F^{(k)})$

- $\mathcal{L}\left(|F^{(k)}| \mid (m_0, \dots, m_{k-1})\right) = \mathcal{L}\left(|\vec{F}^{(k)}| \mid (\sum m_i, 0, \dots, 0)\right)$

- As a process in  $k$ :

- $\mathcal{L}(N^{(k)}, k \geq 0)$

- $\mathcal{L}(\{|O^{(k)}|\}, k \geq 0)$

# Summary of universality results

## Theorem

For any continuous model with valid spreading policy, the following distributions are **explicit** and **independent of the spreading policy**:

- With  $k$  fixed:
  - $\mathcal{L}(O^{(k)}, F^{(k)})$
  - $\mathcal{L}\left(|F^{(k)}| \mid (m_0, \dots, m_{k-1})\right) = \mathcal{L}\left(|\vec{F}^{(k)}| \mid (\sum m_i, 0, \dots, 0)\right)$
- As a process in  $k$ :
  - $\mathcal{L}(N^{(k)}, k \geq 0)$
  - $\mathcal{L}(\{|O^{(k)}|\}, k \geq 0)$

**Corollary:** results on  $O^{(k)}, F^{(k)}$  for one spreading policy are valid for any spreading policy !

With  $n$  (random) masses,  $n \rightarrow \infty$ , for example if

- $\forall i, m_i = 1/n$  and consider the process at time  $n$
- $\forall i, m_i = \ell_i/n$  (where  $\ell_i$  are i.i.d. with  $\mathbb{E}[\ell_i] < \infty$  and satisfy some regularity assumption), at time  $t = \sup\{k : \sum_{i=0}^k m_i < 1\}$ .



With  $n$  (random) masses,  $n \rightarrow \infty$ , for example if

- $\forall i, m_i = 1/n$  and consider the process at time  $n$
- $\forall i, m_i = \ell_i/n$  (where  $\ell_i$  are i.i.d. with  $\mathbb{E}[\ell_i] < \infty$  and satisfy some regularity assumption), at time  $t = \sup\{k : \sum_{i=0}^k m_i < 1\}$ .

## Corollary (Bertoin, Miermont [BM06]; Marckert, V.)

*There exists a limit process  $S$  such that*

$$\left( \frac{\text{LargestBlock}^{(i)}}{n}, 1 \leq i \leq j \right) \xrightarrow[n \rightarrow \infty]{(d)} (\text{SortedExc}(S)_i, 1 \leq i \leq j).$$

# Asymptotic results

With  $n$  (random) masses,  $n \rightarrow \infty$ , for example if

- $\forall i, m_i = 1/n$  and consider the process at time  $n$
- $\forall i, m_i = \ell_i/n$  (where  $\ell_i$  are i.i.d. with  $\mathbb{E}[\ell_i] < \infty$  and satisfy some regularity assumption), at time  $t = \sup\{k : \sum_{i=0}^k m_i < 1\}$ .

## Corollary (Bertoin, Miermont [BM06]; Marckert, V.)


*There exists a limit process  $S$  such that*

$$\left( \frac{\text{LargestBlock}^{(i)}}{n}, 1 \leq i \leq j \right) \xrightarrow[n \rightarrow \infty]{(d)} \left( \text{SortedExc}(e^{(\lambda)})_i, 1 \leq i \leq j \right).$$

# Asymptotic results

With  $n$  (random) masses,  $n \rightarrow \infty$ , for example if

- $\forall i, m_i = 1/n$  and consider the process at time  $n$
- $\forall i, m_i = \ell_i/n$  (where  $\ell_i$  are i.i.d. with  $\mathbb{E}[\ell_i] < \infty$  and  $\mathbb{P}[\ell_i > 0] > 0$ ),  
at time  $t = \sup\{k : \sum_{i=0}^k m_i < 1\}$ .



**Corollary (Bertoin, Miermont [BM06]; Marckert, V.)**



*There exists a limit process  $S$  such that*

$$\left( \frac{\text{LargestBlock}^{(i)}}{n}, 1 \leq i \leq j \right) \xrightarrow[n \rightarrow \infty]{(d)} \left( \text{SortedExc}(e^{(\lambda)})_i, 1 \leq i \leq j \right).$$




# Key tools : the “collecting paths”

## Illustration:



# Key tools : the “collecting paths”


Illustration:



Definition:  $S_x = -x + \sum_{j=0}^{k-1} m_j \mathbb{1}_{u_j \leq x}$ ,  $\forall x \in [0, 1]$


# Key tools : the “collecting paths”

Illustration:




Definition:  $S_x = -x + \sum_{j=0}^{k-1} m_j \mathbb{1}_{u_j \leq x}$ ,  $\forall x \in [0, 1]$

Periodic extension  $\bar{S}$ :




# Key tools : the “collecting paths”

## Illustration:



**Definition:**  $S_x = -x + \sum_{j=0}^{k-1} m_j \mathbb{1}_{u_j \leq x}$ ,  $\forall x \in [0, 1]$

## Periodic extension $\bar{S}$ :




**Convergence:** when  $\bar{S}$  converges to  $e^{(\lambda)}$ ,  
 $(\text{SortedExc}(\bar{S}))_{1 \leq i \leq j} \xrightarrow[n \rightarrow \infty]{(d)}$   $(\text{SortedExc}(e^{(\lambda)}))_{1 \leq i \leq j}$ .

# Discrete space and discrete masses

**New definition** on  $\mathcal{C}_n := \{0/n, \dots, (n-1)/n\} \subset \mathcal{C}$  :

- masses arrive on  $\mathcal{C}_n$  :  $\forall i, u_i \sim \mathcal{U}(\mathcal{C}_n)$
- they cover intervals with extremities in  $\mathcal{C}_n$
- their spreading policy is only invariant by  $1/n$  rotation





# Discrete space and discrete masses

**New definition** on  $\mathcal{C}_n := \{0/n, \dots, (n-1)/n\} \subset \mathcal{C}$  :

- masses arrive on  $\mathcal{C}_n$  :  $\forall i, u_i \sim \mathcal{U}(\mathcal{C}_n)$
- they cover intervals with extremities in  $\mathcal{C}_n$
- their spreading policy is only invariant by  $1/n$  rotation



## Theorem (Similar universality result)

For any **discrete** model with valid spreading policy, the following distributions are **explicit** and **independent of the spreading policy**:

- With  $k$  fixed:
  - $\mathcal{L}(O^{(k)}, F^{(k)})$
  - $\mathcal{L}\left(|F^{(k)}| \mid (m_0, \dots, m_{k-1})\right) = \mathcal{L}\left(|\overrightarrow{F^{(k)}}| \mid (\sum m_i, 0, \dots, 0)\right)$
- As a process in  $k$ :
  - $\mathcal{L}(N^{(k)}, k \geq 0)$
  - $\mathcal{L}(\{|O^{(k)}|\}, k \geq 0)$


# Discrete VS Continuous model

Continuous process:



$$N^{(k)} \stackrel{(d)}{=} 1 + \text{Binomial}(k - 1, R)$$


Discrete process:



$$N^{(k)} \stackrel{(d)}{=} 1 + \dots$$


# Discrete VS Continuous model

Continuous process:



$$N^{(k)} \stackrel{(d)}{=} 1 + \text{Binomial}(k - 1, R)$$

Discrete process:



$$N^{(k)} \stackrel{(d)}{=} 1 + \text{Binomial}(k - 1, R - 1/n)$$


# Discrete VS Continuous model

Continuous process:



$$N^{(k)} \stackrel{(d)}{=} 1 + \text{Binomial}(k - 1, R)$$


Discrete process:



$$N^{(k)} \stackrel{(d)}{=} 1 + \text{NbDiff}(\text{Binomial}(k - 1, R - 1/n))$$


# Discrete VS Continuous model

Continuous process:



$$N^{(k)} \stackrel{(d)}{=} 1 + \text{Binomial}(k-1, R)$$

Discrete process:



$$N^{(k)} \stackrel{(d)}{=} 1 + \text{NbDiff}(\text{Binomial}(k-1, R - 1/n))$$


**Asymtotic behavior** when  $k = n - \lambda\sqrt{n}$  and  $\forall i, m_i = 1/n$

Number of blocks:  $\frac{N_k}{\sqrt{n}} \xrightarrow[n \rightarrow \infty]{\mathbb{P}} \lambda$

Number of blocks:  $\frac{N_k^{(n)}}{\sqrt{n}} \xrightarrow[n \rightarrow \infty]{\mathbb{P}} \lambda(1 - e^{-1})$


# Discrete VS Continuous model

Continuous process:



$$N^{(k)} \stackrel{(d)}{=} 1 + \text{Binomial}(k-1, R)$$

Discrete process:



$$N^{(k)} \stackrel{(d)}{=} 1 + \text{NbDiff}(\text{Binomial}(k-1, R-1/n))$$

**Asymtotic behavior** when  $k = n - \lambda\sqrt{n}$  and  $\forall i, m_i = 1/n$

**Number of blocks:**  $\frac{N_k}{\sqrt{n}} \xrightarrow[n \rightarrow \infty]{\mathbb{P}} \lambda$

**Number of blocks:**  $\frac{N_k^{(n)}}{\sqrt{n}} \xrightarrow[n \rightarrow \infty]{\mathbb{P}} \lambda(1 - e^{-1})$

**Large block sizes:**  $\left( \frac{\text{LargestBlock}^{(i)}}{n}, 1 \leq i \leq j \right) \xrightarrow[n \rightarrow \infty]{(d)} (\text{SortedExc}(e^\lambda))_i, 1 \leq i \leq j$

## Cost of parking procedures:

- Standard parking : if car  $k$  falls in a block of size  $B_k$ , it costs  $C_k = \lfloor U.B_k \rfloor$ , with  $U \sim \mathcal{U}([0, 1])$ . Chassaing-Louchard [CL02]:







$$\frac{1}{n^{3/2}} \sum_{k=1}^{\lfloor n - \lambda\sqrt{n} \rfloor} C_k \xrightarrow[n \rightarrow \infty]{(d)} \int_0^1 e_t^{(\lambda)} dt$$

- Parking with different parking policies : see Marckert-V. for some of them.

💧 Thank you ! 💧



## References

-  David Aldous and Jim Pitman, [The standard additive coalescent](#), *The Annals of Probability* **26** (1998), no. 4, 1703 – 1726.
-  Jean Bertoin and Grégory Miermont, [Asymptotics in Knuth's parking problem for caravans](#), *Random Structures and Algorithms* **29** (2006), no. 1, 38–55, 18 pages, 2 figures.
-  Philippe Chassaing and Guy Louchard, [Phase transition for parking blocks, brownian excursion and coalescence](#), *Random Structures & Algorithms* **21** (2002), no. 1, 76–119.
-  Donald E. Knuth, [The art of computer programming. Volume 3](#), Addison-Wesley Series in Computer Science and Information Processing, Addison-Wesley Publishing Co., Reading, Mass.-London-Don Mills, Ont., 1973, Sorting and searching. MR 445948
-  Alan G. Konheim and Benjamin Weiss, [An occupancy discipline and applications](#), *SIAM Journal on Applied Mathematics* **14** (1966), no. 6, 1266–1274.
-  Michał Przykucki, Alexander Roberts, and Alex Scott, [Parking on the integers](#), *Ann. Appl. Probab.* **33** (2023), no. 2, 876–901. MR 4564421